



PL
RB-54
United States
Department of
Agriculture

Forest
Service


North Central
Forest Experiment
Station

Resource
Bulletin NC-88



Plant Species Biomass Estimates for 13 Upland Community Types of Northeastern Minnesota

Lewis F. Ohmann and David F. Grigal



**North Central Forest Experiment Station
Forest Service—U.S. Department of Agriculture
1992 Folwell Avenue
St. Paul, Minnesota 55108**

**Manuscript approved for publication October 24, 1985
1985**

PLANT SPECIES BIOMASS ESTIMATES FOR 13 COMMUNITY TYPES OF NORTHEASTERN MINNESOTA

Lewis F. Ohmann, *Principal Plant Ecologist,*

Grand Rapids, Minnesota,

and David F. Grigal, Professor,

Departments of Soil Science and Forest Resources,

University of Minnesota,

St. Paul, Minnesota

A series of interrelated studies designed to classify, describe, and elucidate the dynamics of upland plant community types found in northeastern Minnesota has resulted in a unique phytosociological data base. The data base contains detailed information by species and vegetative strata for 13 community types—12 forest and one nonforest. The concurrent development of biomass-estimation equations for most of the species represented in the data set allowed conversion of these phytosociologic data into estimates of biomass for various stages of community development ranging from 1 year following disturbance to maturity.

THE DATA BASE

Field

The data base represented in the accompanying tables is the result of phytosociologic and biomass data collected within or near the Boundary Waters Canoe Area Wilderness (BWCAW) as part of studies of: (1) mature upland forest and rock outcrop stands that regenerated naturally following disturbance, usually wildfire (Ohmann and Ream 1971a); (2) mature upland forest stands that regenerated naturally following logging of red and white pine, usually followed by slash fires, during the 1890 to 1930 era (Grigal and Ohmann 1975); (3) 33-year-old upland forest stands regenerated naturally following summer-season wildfires in 1936 (Ohmann *et al.* 1973); (4) 1- to 30-year-old upland forest stands regenerated by planting following logging of jack pine and black spruce (Ohmann 1982, 1984); (5) 1- to 5-year-old upland forest stands that regenerated naturally following a 1971 spring-season fire (Ohmann and Grigal 1979); and (6) 1-year-old forest stands that regenerated naturally following a 1974 summer-season wildfire (Ohmann and Grigal 1981) (table 1).¹

The sampling design for studies 1 through 4 has been reported in detail (Ohmann and Ream 1971b). Briefly, we randomly located stands by using a grid overlay on standard topographic maps. Only upland plant communities—defined as those on which surface water never accumulates—were sampled. The minimum size of a sample area was about 5 acres (2 ha) in communities with trees and about 1/40 acre (100 m²) in communities without trees (i.e., bedrock outcrops). Only uniform sites, with no apparent discontinuities in either vegetation or topography, were selected. We used 20 points within each stand to sample the vegetation. The point-centered quarter method (Cottam and Curtis 1956) was used for trees; circular milacre plots (4 m²) were used for tree seedlings (<1.0 in. (2.5 cm) d.b.h.) and tall shrubs; and 1 × 2-ft (0.2-m²) rectangular plots were used for other vegetation. The sample design and vegetation descriptions for studies 5 and 6 have been reported in detail (Ohmann and Grigal 1979, 1981). Biomass values were developed directly from vegetation clipped from 6.5-ft² (0.6-m²) plots established each year adjacent to permanent plots used to sample vegetation for phytosociological characteristics.

Numerical

Trees and saplings

For those studies of forest stands old enough to have saplings (1- to 4-in. (2.5 to 10 cm) d.b.h.) and trees (>4 in. (10 cm) d.b.h.), biomass estimation equations were based on the abundant recent literature (table 2). Aboveground total and foliage biomass were estimated separately for each species and summarized for each community type. For uniformity and consistency, we used the same functional relationship for all tree and sapling biomass

¹All tables appear at the end of the text.

estimation equations. Because of its wide use, we selected the allometric relationship

$$y = aD^b , \quad (1)$$

where y is mass of a tree component, D is diameter at breast height (d.b.h.), and a and b are regression parameters. When available, general equations from many sites were used (e.g., Crow 1983, Schmitt and Grigal 1981). In some cases our search only indicated one set of equations for a species (e.g., black ash, (Parker and Schneider 1975)). In other cases, several equations, representing a wide geographic range, were available for a species. Regression coefficients were generally similar for a species, independent of location. We pooled the parameters for equations that were similar to arrive at a more generalized expression. The approach was similar to that used by Pastor *et al.* (1984).

The usual method of computation for point-quarter data uses stem diameter and distance from a number of points (usually 20 or more). The data are then summed over all points, and tree density and basal area are calculated on the basis of the aggregated data (Cottam and Curtis 1956). Our approach was to perform the appropriate computations for data from each point. Summary statistics, including mean and standard error (SE), could then be developed over all points. The total biomass for a stratum was the sum of the biomass for each species in the stratum.

For studies 3 and 4, all tree stems greater than 1 in. (2.5 cm) were combined; thus, there is no separate biomass estimate for saplings. In studies 5 and 6, all tree stems were classified as tree seedlings (table 1).

Tall shrubs

Biomass estimation equations for tall shrubs were also of the allometric form. They were based primarily on three studies from northern Minnesota: Balogh (1983), Grigal and Ohmann (1977), and Roussopoulos and Loomis (1979) (table 3). Equations from Telfer (1969) were used for a few less common species. For some common species, all three primary sources contained estimation equations. In those cases, all three equations were plotted and the intermediate equation was selected. Because this was often an equation from Balogh (1983), equations for less common species were also obtained from that source. All equations were based on shrub diameter at 6 in. (15 cm) above the ground.

In studies 1 through 3, tall shrubs were measured in 0.4-in. (1-cm) diameter classes at 6 in. (15 cm) above the ground (Ohmann and Ream 1971a) (table 1). However, other data (Balogh 1983) indicate that the majority of shrubs in smaller size classes are near the lower end of these classes. Use of the allo-

metric function based on 0.4-in. (1-cm) diameter classes would therefore overestimate shrub biomass. We determined diameter-density distributions for each shrub species in studies 1 through 3. The data were fit to a linearized power function

$$\ln (De) = a + b \ln (Di) , \quad (2)$$

where $\ln (De)$ is the natural log of proportion of total stems of that diameter, and $\ln (Di)$ is the natural log of stem diameter. The slope of this equation (b) for each species is the change in proportion of individuals with diameter. Shrub data for a sampling point, in 0.4-in. (1-cm) diameter classes, and the slope (b) of the above relationship were used to compute the expected number of individuals by 0.1-in. (0.25-cm) diameter classes. In study 4 the number of shrub stems by 0.1-in. (0.25-cm) diameter classes were determined directly within each plot (table 1). The biomass estimation equations were then applied to the 0.1-in. (0.25-cm) classes. Biomass was calculated for each tall shrub species on a plot. The means of those plot values were calculated by species and summed for total biomass for the shrub stratum. In studies 5 and 6, the aboveground standing crop of vegetation was sampled during August of each year following fire (only the first year in the case of the summer-season fire, study 6). All shrub vegetation in 10 (20 in study 6) randomly located plots near the permanent sample plots within each stand was clipped at ground level (table 1). All harvested shrub vegetation was returned to the laboratory, sorted, counted, and bagged by species and plot, dried to 70° C to constant weight, and weighed. Means and variances of the samples were calculated for all species in each stand.

Tree seedlings

Tree seedling cover and density were recorded in studies 1 through 4 (table 1). Most published biomass estimation equations for seedlings are based on either diameter (Telfer 1969, Roussopoulos and Loomis 1979) or height (Young *et al.* 1980). We modified the published equations to determine biomass for seedlings from our data.

Our independent variable was cover per individual seedling. We collected samples to determine minimum and maximum biomass and cover for representative deciduous (northern red oak, red maple, quaking aspen, and paper birch) and coniferous (balsam fir, black spruce, jack pine, and white pine) species that were most common as seedlings in the data set.

Relationships from both Roussopoulos and Loomis (1979) and Young *et al.* (1980) were used to determine change in mass with size. Biomass estimation equations of the allometric form were then developed for both foliage mass and for total mass, using

cover per individual as the independent variable. In studies 1 through 4, seedling biomass for a stand was calculated similarly to that for tall shrubs. In studies 5 and 6, biomass was determined by clipping and weighing as for tall shrubs (table 1).

Herbs and low shrubs

A series of biomass equations using cover as the independent variable has been developed for herbs, ferns, and low shrubs sampled on $1 \times 2\text{-ft}$ (0.2-m^2) plots (Ohmann *et al.* 1981) (table 1). These equations were used for the data from studies 1 through 4. For species for which no equation had been specifically developed, we used equations for similar species (e.g., *Fragaria* spp. equation for *Waldsteinia fragarioides* (Michx.) Tratt.). A miscellaneous herb and fern equation (unpublished) was used for species with no obvious morphological match. For each species, stand biomass was simply the mean of the estimates for each plot (usually 20), and standard error was calculated. Stand totals over all species were calculated as with the other strata. Biomass for studies 5 and 6 was obtained by clipping plots exactly as for shrubs and tree seedlings (table 1).

Mosses and lichens

Cover estimates for both mosses and lichens were also made on $1 \times 2\text{-ft}$ (0.2-m^2) plots in studies 1 through 3 (table 1). Based on sampling in northeastern Minnesota, we determined mass per unit area for various moss and lichen species and species groups (Ohmann and Grigal, 1985). These data were then converted to linear relationships between cover and mass on $1 \times 2\text{-ft}$ (0.2-m^2) plots. Means and errors were calculated as for the herb stratum. The values obtained are within the range of those reported for jack pine stands on shallow soils in the area (Green and Grigal 1979). Moss and lichen cover was not estimated in study 4 (table 1). In studies 5 and 6, moss and lichen biomasses were determined from the clip plots exactly like those for the other vegetation (table 1).

Each immature stand comprising studies 2 through 6 was assigned to one of 12 forest community types using a classification technique applied earlier to stands in studies 1 and 2 (Grigal and Ohmann 1975). The assignment was made by multiplying the frequency of occurrence of each of the 53 indicator species present in a stand by a classification function for that species (table 4). The sum of those products together with the appropriate constant (table 4) indicated a stand's affinity to each of the 12 community types. Stands were assigned to the community type for which they showed the greatest affinity. The assignment of stands only 1 year from fire or logging disturbance, as well as those 33 years from disturbance was possible because the majority

of the 53 indicator species are neither tree nor disturbance-related species.

Tables 5-17 present the estimated aboveground ovendry biomass by species and stratum for each of the 13 community types. Means and standard errors are presented except where a species occurred in only one stand of a community type.

Thirteen community types were recognized in the initial floristic classification (Grigal and Ohmann 1975). The *lichen* community type consists of lichen-covered bedrock outcrops on ridgetops and upper slopes. Although overstory trees are absent, the type contains a few woody seedlings and tall shrubs, and some low shrubs and herbs. The remaining 12 types are forested. The *jack pine-oak* community type occurs in similar physiographic situations as the lichen type, except that sufficient soil is present to support a forest community. The community is dominated by jack pine, with some northern red oak and red maple. Where these stands have been severely disturbed by logging and fire, the coniferous component has been markedly reduced, giving rise to the *maple-oak* community type. Red maple and northern red oak dominate the overstory in this type, but the understory is similar to that in the jack pine-oak type.

Two other community types are dominated by jack pine. The *jack pine-fir* type is on deep soils, and has an overstory of jack pine with balsam fir, paper birch, and black spruce saplings and seedlings. A tall shrub layer is common, and mountain maple and fly honeysuckle are important. Large-leaf aster dominates the herb layer, but a variety of other herbs occur in the type. The *jack pine-black spruce* type, as the name implies, contains an overstory of jack pine with black spruce trees, saplings, and seedlings. Tall shrubs are rare, and moss species rather than herbaceous species dominate the forest floor. In the *black spruce-feather moss* community type, black spruce dominates the overstory although some jack pine is present. Shrubs and herbs are less common than in the previous type, and the moss layer covers the entire forest floor.

Four types are partially to completely dominated by quaking aspen and paper birch. The *aspen-birch* community type is dominated by those two species, although balsam fir seedlings are abundant. Tall shrubs are plentiful, dominated by beaked hazel. Low shrubs, especially bush honeysuckle are also abundant. Large-leaf aster is an important herb species, along with false lily-of-the-valley, wild sarsaparilla, and bunchberry. Mosses are unimportant. The *aspen-birch-white pine* community type is similar to the aspen-birch type, but it contains a component of large white pine and a greater abundance of

balsam fir in the overstory. The *maple-aspen-birch* community type contains red maple, especially as a seedling, in addition to quaking aspen and paper birch. Tall shrubs include green alder and mountain maple, in addition to beaked hazel. The type also contains balsam fir, but that species is more important in the *maple-aspen-birch-fir* community type. The type is similar to the *maple-aspen-birch* type, but mountain maple is more dominant as a tall shrub, low shrubs are less frequent, and bunchberry is the most common herb.

Both the conifer and the broadleaf types previously mentioned appear to occur in a successional sequence to the *fir-birch* community type. Balsam fir is the most common tree, with both paper birch and northern white-cedar as associates. Mountain maple is the dominant tall shrub, and low shrubs are sparse. The herb layer is much reduced compared to the broadleaf types, with bunchberry the most abundant herb. With the decline of the herb stratum, mosses are more abundant. The *white-cedar* community type appears to be even further along the successional sequence than the *fir-birch* type, and the stands in this type are somewhat older than those in the previously mentioned types. The type is similar to the *fir-birch* type, but northern white-cedar is more important and the herb stratum is more mixed. Bishop's cap is a common herb.

The final type, *red pine* does not fit well in sequence with the other types. It consists of old-growth red pine with a poorly developed understory resulting in open, nearly park-like stands. White pine is a common associated species. Low ericaceous shrubs are common, as are mosses and lichens.

The validity of our results is predicated in part on the generality of biomass estimation equations. Such generality across sites and even across regions has been examined for trees, and relations between diameter and total aboveground biomass are very consistent for a number of species included in our data set. These species include red maple (Crow 1983), paper birch (Schmitt and Grigal 1981), jack pine (Green and Grigal 1978), black spruce (Grigal and Kernik 1984), red pine (Alban and Laidly 1982), and quaking aspen, white pine, and northern red oak (Pastor *et al.* 1984). These results lend credence to our approach. Extensive tests of generality have not been conducted for most species in the lower forest strata, but in this study most of those equations were derived from data collected in or near the BWCAW.

Biomass productivity and the dynamic biomass relations among and within community types represented by the estimates reported here, along with their wildlife implications, have been discussed elsewhere (Grigal *et al.* 1976; Grigal and Ohmann 1977;

Ohmann *et al.* 1976, 1978; Ohmann and Grigal 1977, 1979, 1981, accepted; Ohmann 1982, 1984).

LITERATURE CITED

- Alban, D. H.; Laidly, P. R. Generalized biomass equations for jack and red pine in the Lake States. *Can. J. For. Res.* 12: 913-921; 1982.
- Balogh, J. C. Tall shrubs in northern Minnesota. St. Paul, MN: University of Minnesota Soil Science Department; 1983. 140 p. Ph.D. thesis.
- Barney, Richard J.; Van Cleve, Keith; Schlentner, Robert. Biomass distribution and crown characteristics in two Alaskan *Picea mariana* ecosystems. *Can. J. For. Res.* 8 (1): 36-41; 1978.
- Baskerville, G. L. Dry matter production in immature balsam fir stands. *For. Sci. Monogr.* 9: 42 p; 1965.
- Bella, I. E.; De Franceschi, J. P. Biomass productivity of young aspen stands in western Canada. Inf. Rep. X-219. Edmonton, Alberta: Canadian Forestry Service, Northern Forest Research Centre; 1980: 1-13.
- Cottam, G.; Curtis, J. T. The use of distance measures in phytosociological sampling. *Ecology* 37: 451-460; 1956.
- Crow, T. R. Comparing biomass regressions by site and stand age for red maple. *Can. J. For. Res.* 13: 283-288; 1983.
- Green, D. E.; Grigal, D. F. Jack pine biomass accretion on shallow and deep soils in Minnesota. *Soil Sci. Soc. Am. J.* 43: 1233-1237; 1979.
- Green, D. E.; Grigal, D. F. Generalized biomass estimation equations for jack pine. MN For. Res. Notes 268. St. Paul, MN: College of Forestry; 1978. 4 p.
- Grigal, D. F.; Kernik, L. K. Biomass estimation equations for black spruce (*Picea mariana* (Mill.) B.S.P.); For. Res. Note 290. St. Paul, MN: University of Minnesota College of Forestry, 1984. 4 p.
- Grigal, D. F.; Ohmann, L. F. Classification, description and dynamics of upland communities within a Minnesota wilderness area. *Ecol. Monogr.* 45: 389-407; 1975.
- Grigal, D. F.; Ohmann, L. F. Biomass estimation for some shrubs from northeastern Minnesota. Res. Note NC-226. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1977. 3 p.
- Grigal, D. F.; Ohmann, L. F.; Brander, R. B. Seasonal dynamics of tall shrubs in northeastern Minnesota: biomass and nutrient element change. *For. Sci.* 22(2): 195-208; 1976.
- Harding, R. B. Site quality evaluation for white spruce plantations in northern Minnesota. St. Paul, MN: University of Minnesota College of Forestry; 1982. 135 p. Ph.D. thesis.

- Ker, M. F. Tree biomass equations for ten major species in Cumberland, Nova Scotia. Maritime For. Res. Cent. M-X-108. Fredericton, New Brunswick, Canada: Maritime Forest Research Centre; 1980. 26 p.
- Kinerson, A. S.; Bartholomew, I. Biomass estimation equations and nutrient composition of white pine, white birch, red maple, and red oak in New Hampshire. Res. Rep. 62. Durham, NH: Agriculture Experiment Station; 1977. 8 p.
- Koerper, Greg L.; Richardson, Curtis J. Biomass and net annual production regressions for *Populus grandidentata* on three sites in northern lower Michigan. Can. J. For. Res. 10: 92-101; 1980.
- Monteith, D. B. Whole tree weight tables for New York. Res. Rep. 40. Syracuse, NY: State University of New York, Applied Forest Research Institute; 1979. 64 p.
- Ohmann, Lewis F. Tall shrub layer biomass in conifer plantations of northeastern Minnesota. Res. Pap. NC-219. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1982. 30 p.
- Ohmann, Lewis F. Biomass in conifer plantations of northeastern Minnesota. Res. Pap. NC-247. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1984. 25 p.
- Ohmann, L. F.; Grigal, D. F. Some individual plant biomass values from northeastern Minnesota. Res. Note NC-227. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1977. 2 p.
- Ohmann, Lewis F.; Grigal, D. F. Early revegetation and nutrient dynamics following the 1971 Little Sioux Forest Fire in northeastern Minnesota. For. Sci. Monogr. 21; 1979. 80 p.
- Ohmann, L. F.; Grigal, D. F. Contrasting vegetative responses following two forest fires in northeastern Minnesota. Am. Midl. Nat. 106(1): 54-64; 1981.
- Ohmann, Lewis F.; Grigal, D. F. Biomass distribution of unmanaged upland forests in Minnesota. For. Ecol. and Manage. 13: 1985.
- Ohmann, Lewis F.; Ream, Robert R. Wilderness ecology: virgin plant communities of the Boundary Waters Canoe Area. Res. Pap. NC-63. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1971a. 55 p.
- Ohmann, Lewis F.; Ream, Robert R. Wilderness ecology: method of sampling and summarizing data for plant community classification. Res. Pap. NC-49. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1971b. 14 p.
- Ohmann, L. F.; Grigal, D. F.; Brander, R. B. Biomass estimation for five shrubs from northeastern Minnesota. Res. Pap. NC-133. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1976. 11 p.
- Ohmann, L. F.; Grigal, D. F.; Brander, R. B. Shrub mass: data variability and changes over time. In: Proceedings, 10th North American Moose Workshop; Duluth, MN: 172-214; 1978.
- Ohmann, Lewis F.; Grigal, David F.; Rogers, Lynn L. Estimating plant biomass for undergrowth species of northeastern Minnesota communities. Gen. Tech. Rep. NC-61. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1981. 10 p.
- Ohmann, L. F.; Cushwa, C. T.; Lake, R. E.; Beer, J. R.; Brander, R. B. Wilderness ecology: the upland plant communities, woody browse production, and small mammals of two adjacent 33-year-old wildfire areas of northeastern Minnesota. Gen. Tech. Rep. NC-7. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1973. 30 p.
- Parker, G. R.; Schneider, G. Biomass and productivity of an alder swamp in northern Michigan. Can. J. For. Res. 5: 403-409; 1975.
- Pastor, John P.; Bockheim, J. G. Biomass and production of an aspen-mixed hardwood-spodosol ecosystem in northern Wisconsin. Can. J. For. Res. 11: 132-138; 1981.
- Pastor, J.; Aber, J. D.; Meililo, J. M. Biomass prediction using generalized allometric regressions for some Northeast tree species. For. Ecol. and Manage. 7: 265-274; 1984.
- Pollard, D. F. W. Above-ground dry matter production in three stands of trembling aspen. Can. J. For. Res. 2: 27-33; 1972.
- Roussopoulos, P. J.; Loomis, R. M. Weights and dimensional properties of shrubs and small trees of the Great Lakes conifer forests. Res. Pap. NC-178. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1979. 6 p.
- Schmitt, Mark D. C.; Grigal, D. F. Generalized biomass estimation equations for *Betula papyrifera* Marsh. Can. J. For. Res. 11(4): 837-840; 1981.
- Telfer, E. S. Weight-diameter relationships for 22 woody plant species. Can. Bot. 47: 1851-1855; 1969.
- Whittaker, R. H.; Woodwell, G. M. Dimension and production relations of trees and shrubs in the Brookhaven Forest, New York. J. Ecol. 56: 1-25; 1968.
- Young, Harold E.; Ribe, John H.; Wainwright, Kevin. Weight tables for tree and shrub species in Maine. Misc. Rep. 230. Orono, ME: University of Maine Life Sciences and Agriculture Experiment Station; 1980. 84 p.

Table 1.--Type and source of data used to develop biomass estimates for species and stratum for 13 plant community types of northeastern Minnesota

	Prayer Lake Fire	Little Sioux Fire	Isabella Plantations	Cherokee- Frost Lake	BWCAW ^{1/}
Age (years)	1	1-5	1-30	33	mature
Number of stands sampled	2	7	53	35	121
Tree foliage & total above- ground biomass	None	None	Stems >1 in. d.b.h. on 20 plotless points	Trees >4 in. d.b.h. Saplings 1-4 in. d.b.h. on 20 plotless points	
Tree seedling foliage & total aboveground biomass	Stems clipped in 20 6.5-ft ² plots	Stems clipped in 10 6.5-ft ² plots	Percent cover of seedlings <1 in. d.b.h. on 20 acre plots		
Tall shrub foliage & total aboveground biomass	Stems clipped in 20 6.5-ft ² plots	Stems clipped in 10 6.5-ft ² plots	Stems tallied in 0.1 in. diameter classes at 6 in. aboveground in 10 acre plots	Stems tallied in 0.4 in. diameter classes at 6 in. aboveground in 20 acre plots	
Herb (foliage) & low shrub (total) biomass	Stems clipped in 20 6.5-ft ² plots	Stems clipped in 10 6.5-ft ² plots	Species percent ground cover on 20 2-ft ² plots		
Moss & lichen foliage biomass	Clipped in 20 6.5 ft ² plots	Clipped in 10 6.5- ft ² plots	Not collected	Species percent ground cover on 20 2-ft ² plots	

1/ Boundary Waters Canoe Area Wilderness

Table 2.--Source of biomass estimation equations used for tree and sapling strata

Species	Source
<u>Abies balsamea</u>	Baskerville (1965) Ker (1980) Young et al. (1980)
<u>Acer rubrum</u>	Crow (1983)
<u>Betula papyrifera</u>	Schmitt and Grigal (1981)
<u>Fraxinus nigra</u>	Parker and Schneider (1975)
<u>Larix laricina</u>	Young et al. (1980)
<u>Picea glauca</u>	Harding (1982) Ker (1980) Young et al. (1980)
<u>Pinus banksiana</u>	Green and Grigal (1978)
<u>Picea mariana</u>	Barney et al. (1978) Grigal and Kernik (1983) Ker (1980)
<u>Pinus resinosa</u>	Alban (personal communication 1982) Ker (1980) Young et al. (1980)
<u>Pinus strobus</u>	Kinerson and Bartholomew (1977) Monteith (1979) Young et al. (1980)
<u>Populus balsamifera</u>	Bella and De Franceschi (1980)
<u>Populus grandidentata</u>	Koerper and Richardson (1980)
<u>Populus tremuloides</u>	Ker (1980) Pastor and Bockheim (1981) Pollard (1972)
<u>Prunus pensylvanica</u>	Young et al. (1980)
<u>Quercus rubra</u>	Monteith (1979) Whittaker and Woodwell (1968)
<u>Sorbus americana</u> (<u>Prunus pensylvanica</u> eq.)	Young et al. (1980)
<u>Thuja occidentalis</u>	Young et al. (1980)
<u>Tilia americana</u>	Koerper and Richardson (1980)

Table 3.--Source of biomass estimation equations used for tall shrub data

Species	Source
<u>Acer spicatum</u>	Balogh (1983)
<u>Alnus crispa</u>	Balogh (1983)
<u>Alnus rugosa</u>	Grigal and Ohmann (1977)
<u>Amelanchier</u> spp.	Balogh (1983)
<u>Comptonia peregrina</u>	Telfer (1969)
<u>Cornus rugosa</u>	Balogh (1983)
<u>Cornus stolonifera</u> (<i>C. alternifolia</i> eq.)	Balogh (1983)
<u>Corylus cornuta</u>	Balogh (1983)
<u>Diervilla lonicera</u>	Roussopoulos and Loomis (1979)
<u>Juniperus communis</u> (<i>Abies balsamea</i> eq.)	Roussopoulos and Loomis (1979)
<u>Ledum groenlandicum</u>	Telfer (1969)
<u>Lonicera canadensis</u>	Roussopoulos and Loomis (1979)
<u>Lonicera hirsuta</u>	Grigal and Ohmann (1977)
<u>Lonicera oblongifolia</u>	Grigal and Ohmann (1977)
<u>Lonicera villosa</u> (<i>L. hirsuta</i> eq.)	Grigal and Ohmann (1977)
<u>Prunus pensylvanica</u>	Grigal and Ohmann (1977)
<u>Prunus pumila</u> (<i>Lonicera canadensis</i> eq.)	Roussopoulos and Loomis (1979)
<u>Prunus virginiana</u>	Balogh (1983)
<u>Rhus glabra</u> (<i>Sorbus americana</i> eq.)	Roussopoulos and Loomis (1979)
<u>Ribes</u> spp.	Grigal and Ohmann (1977)
<u>Rosa acicularis</u>	Roussopoulos and Loomis (1979)
<u>Salix</u> spp.	Balogh (1983)
<u>Sambucus</u> spp. (<i>Sorbus americana</i> eq.)	Roussopoulos and Loomis (1979)
<u>Sorbus americana</u>	Roussopoulos and Loomis (1979)
<u>Spiraea</u> spp.	Telfer (1969)
<u>Taxus canadensis</u> (<i>Abies balsamea</i> eq.)	Roussopoulos and Loomis (1979)
<u>Viburnum edule</u> (<i>Prunus pensylvanica</i> eq.)	Grigal and Ohmann (1977)
<u>Viburnum rafinesquianum</u>	Balogh (1983)

Table 4.—Classification functions to assign stands into appropriate community types identified in northeastern Minnesota

Indicator species	Aspen-birch-white pine	Aspen-birch	Fir-birch	Maple-aspen-birch	Maple-aspen-fir-birch	Maple-oak	Jack pine-oak	Red pine	Jack pine-fir	Jack pine-black spruce	Black spruce-feather moss	White-cedar
<i>Calliergonella schreberi</i>	-0.0903	-0.3780	0.4420	-0.3151	-0.1918	0.7837	-0.1613	0.3761	0.5763	0.8901	0.6096	0.5268
<i>Cladonia rangiferina</i>	-1.135	-34.74	1.928	-1.4610	-1.7965	1.0937	-14.77	.5622	.2689	4.1010	.2332	.0972
<i>Cladonia mitis</i>	1.9893	1.7888	1.9739	1.7460	1.7808	5.2556	1.0476	3.0162	1.4204	2.0224	2.1997	2.3009
<i>Dicranum spp.</i>	.4911	.6014	.0936	.6518	.5317	.2368	.3632	.5081	.0822	.0415	.1361	.0061
<i>Polytrichum spp.</i>	-1.437	.0408	-3.713	-6.958	-6.645	-1.1193	-2.3929	-3.610	-0.0730	-3.0583	-2.431	-7.975
<i>Lycopodium clavatum</i>	.2853	.5116	-5.188	.5444	.5525	-1.9534	.1986	.1467	.4872	.7065	.4886	-1.0251
<i>Lycopodium obscurum</i>	-2.733	-2.554	.0046	.6757	.5490	1.1345	-7.151	.0516	.2200	.3213	.2377	.2515
<i>Pteridium aquilinum</i>	1.1561	2.0336	.6038	1.2004	.8902	-1.636	.1076	-1.2221	.0351	.4114	.1180	.2076
<i>Aralia nudicaulis</i>	.5007	.4640	.4214	.6571	.5342	-4.128	.4383	.0346	.0948	.0386	.0906	.5345
<i>Aster macrophyllus</i>	.3937	.3140	.3009	.2752	.1454	.1637	.1270	.1210	.2065	.2273	.2680	
<i>Clintonia borealis</i>	.1203	.1635	-.2513	.2787	.1466	.4605	.2806	.0417	.2182	.4093	.3847	.4070
<i>Cornus canadensis</i>	-.6169	-.6169	.0637	-.7038	-.4629	-.4501	-.4111	-.6665	.0449	.0603	-.1008	.4449
<i>Fragaria vesca</i>	.2479	.5703	-.1717	.1779	-.0076	-.3006	-.0134	-.0585	.2226	.5241	-.3371	.3225
<i>Gaultheria trifolia</i>	1.6988	1.7966	-.3324	1.5242	.8675	-.2394	.7194	1.0174	-.2504	.5315	.3036	1.1082
<i>Limnaea borealis</i>	-.3621	-.2291	.2201	-.5033	-.3464	.2727	-.4541	-.3646	.1625	.1738	.2060	.4318
<i>Maianthemum canadense</i>	.2499	-.0182	-.1599	.2712	.2165	.0724	.3460	.2722	-.0597	.0722	-.0241	-.3332
<i>Melampyrum lineare</i>	1.1665	9.151	-.3369	1.3177	1.0240	.9463	1.2092	.3502	.3633	.4160	.0566	
<i>Nitella nuda</i>	-.3069	-.4460	.7263	-.2426	.0888	.4828	.0429	-.1122	.4573	.5915	.5013	2.0189
<i>Trientalis borealis</i>	-.9979	-.8824	-.2187	-.9076	-.6809	.3780	-.3989	-.4890	-.0833	.0838	.0229	.3994
<i>Viola spp.</i>	-1.3748	-.4491	-.0920	-.4095	-.8465	.4470	-.5823	-.6440	.0837	.2571	.0473	.5383
<i>Streptopus roseus</i>	.8397	.9342	-.0201	.5864	.2956	-.0739	.0112	.6923	-.1136	.0439	.0656	.9285
<i>Gaultheria procumbens</i>	-.4103	-.0873	-.2856	-.9505	-.8432	1.5910	.5420	.1742	-.0706	.0830	-.0059	-.2154
<i>Rubus strigosus</i>	-.1.0433	-.8948	.5142	-.8499	-.3914	.8119	-.4625	-.2654	.4621	.8624	.6421	.8716
<i>Rubus pubescens</i>	-.3224	-.1877	.0386	-.9632	-.9632	.9043	-.7115	.3015	.3147	.2474	.2350	.2965
<i>Vaccinium angustifolium</i>	-.9636	-.1.1873	-.1084	-.9776	-.5946	1.0526	-.3206	.1780	-.0036	.4424	.1920	.2742
<i>Vaccinium myrtilloides</i>	-.5944	-.5888	-.3624	-.5836	-.5765	-.1.8859	-.2530	1.3408	-.5522	.6119	.5282	.2892
<i>Diervilla lonicera</i>	-.3362	.6351	.1233	.2976	.2004	.2402	.0364	.0336	.0336	.1098	.1196	.0456
<i>Abies balsamea (S) /</i>	.3263	.2547	-.2153	.3348	.3325	-.0484	.1863	.0984	-.0488	.0088	.1438	.2333
<i>Acer rubrum (S)</i>	-.0265	-.1992	-.0292	.7124	.7136	-.2496	.5497	.3272	-.1400	.0462	.1130	-.0030
<i>Betula papyrifera (S)</i>	-.0794	.3542	.3245	-.2873	-.0530	.2134	-.1346	-.0976	.2301	.3138	.2903	.6389
<i>Picea glauca (S)</i>	.5360	.2296	-.0702	.1008	-.1386	.2511	-.0216	.5873	.2078	.1991	.2149	
<i>Picea mariana (S)</i>	-.9088	-.9331	-.5845	-.8006	-.6463	.5108	-.2687	-.0243	.2146	.0327	-.1835	
<i>Pinus strobus (S)</i>	.2888	.5700	.1181	.3669	.2153	.5553	-.0017	.0708	.0382	.1969	.1969	.2054
<i>Populus tremuloides (S)</i>	.4541	.6895	-.0240	.2672	.1329	.3265	.0349	.2952	.0136	.1291	.0085	.2381
<i>Thuja occidentalis (S)</i>	-.1160	-.0503	.3086	.0409	.0694	-.1941	-.0611	-.0708	.1416	.0709	.1181	.8337
<i>Quercus rubra (S)</i>	.1265	.0637	.2032	.2732	.1046	-.7440	.3809	-.0155	.0082	.1401	.0777	.1569
<i>Acer spicatum</i>	.1967	.1604	.5146	.5440	.0771	.4430	.3185	.2286	.2885	.3179	.3179	.3642
<i>Alnus crispa</i>	.0987	.4195	-.1157	.5654	.3123	-.4360	.1706	.3763	-.2387	.1158	.1158	.4959
<i>Anelachier spp.</i>	-.5840	-.6236	-.3453	-.7123	-.4157	.4973	-.0711	-.1733	.0696	.1564	.0698	.3236
<i>Cornus rugosa</i>	-.4043	.3280	-.3508	-.6338	-.6781	-.1307	-.4636	-.3303	-.0917	.1249	.1626	.3614
<i>Corylus cornuta</i>	.0022	-.0765	.2353	.0551	.0937	.2555	-.0481	.1723	.1799	.3524	.2847	.3612
<i>Lonicera canadensis</i>	-.6864	-.8248	-.4273	-.7290	-.6125	-.1.990	-.0564	-.5220	-.1.694	.2195	.4369	.2998
<i>Salix spp.</i>	-.1754	-.3397	-.0480	-.3962	-.2092	.2529	-.7527	-.1.0611	.2654	.0766	-.0073	.3905
<i>Populus tremuloides (T)</i>	.7950	1.2483	1.0487	.8616	-.1711	.5677	.6049	.0027	.0457	.1345	.1388	.1579
<i>Betula papyrifera (T)</i>	1.1644	1.2591	.5830	1.1523	.8694	.0645	.4404	.6642	.2776	.1345	.3595	.2971
<i>Abies balsamea (T)</i>	.6302	.6948	.4677	.5584	.5177	.1859	.2920	.3499	.3312	.1713	.2909	.2457
<i>Thuya occidentalis (T)</i>	.3133	.3119	.8577	.3432	.4914	.7283	.1869	.3076	.5547	.6218	.6385	.1.3228
<i>Pinus banksiana (T)</i>	.2012	-.0439	.1315	.2378	.1372	.2302	.2519	.3005	.2519	.4015	.1479	.1592
<i>Pinus taeda (T)</i>	.4138	.5196	-.1152	.6869	.4432	-.8575	.2395	-.0893	.2459	.5936	.3820	.1.7174
<i>Pinus mariana (T)</i>	.3650	.2754	.6489	.3673	.4928	.3494	.2426	.3848	.3844	.6265	.6685	.7407
<i>Pinus resinosa (T)</i>	.6742	.3823	.1344	.7702	.5826	.4093	.5975	1.2845	.2915	.2838	-.0877	
<i>Pinus strobus (T)</i>	1.0096	.7417	.3527	.9743	.3111	.0072	.4813	.9342	.1039	.2039	.3999	.1565
<i>Acer rubrum (T)</i>	1.2837	1.3708	.5701	1.5662	1.2423	-.2180	.8816	.5616	.1884	.0685	.2520	.4052
Constant		-140.9238	-152.5211	-95.9515	-169.9262	-130.0939	-205.5043	-95.6863	-154.3386	-61.7259	-90.2644	-77.3991
												-179.2447

1/(T) Tree size class (>1 in. d.b.h.)
(S) Seeding size class (<1 in. d.b.h.)

Table 5.--Aboveground ovendry biomass by species and stratum for the lichen community type of northeastern Minnesota

Stratum/Species	(In Kg/ha) BWC AW			
	Foliage		Mature (n = 6)	
	\bar{X}	SE	\bar{X}	SE
Tree seedling				
<i>Acer rubrum</i> L.	0.5	-	16.0	-
<i>Betula papyrifera</i> Marsh.	.1	-	2.6	-
<i>Pinus banksiana</i> Lamb.	.6	-	4.6	4.6
<i>Populus tremuloides</i> Michx.	.5	-	2.8	-
<i>Quercus rubra</i> L.	.3	-	1.5	-
<i>Sorbus americana</i> Marsh.	.1	-	.7	-
Stratum Total	2.1		28.2	
Tall shrub				
<i>Acer spicatum</i> Lam.	<.1	-	<.1	-
<i>Amelanchier</i> Medic. spp.	<.1	-	.2	.2
<i>Prunus pensylvanica</i> L.F.	<.1	-	<.1	-
<i>Salix bebbiana</i> Sarg.	<.1	-	.1	-
<i>Salix</i> L. spp.	1.0	-	1.2	-
Stratum total	1.1		1.6	
Low shrub				
<i>Diervilla</i> Lonicera Mill.			54.4	53.4
<i>Rubus strigosus</i> Michx.			3.5	-
<i>Vaccinium angustifolium</i> Ait.			37.9	18.7
Stratum total			95.8	
Herb				
<i>Anemone quinquefolia</i> L.	<.1	-		
<i>Aralia hispida</i> Vent.	1.9	-		
<i>Aralia nudicaulis</i> L.	.2	-		
<i>Aster macrophyllus</i> L.	.3	-		
<i>Carex</i> L. spp.	5.0	2.3		
<i>Corydalis sempervirens</i> (L.) Pers.	.5	.3		
<i>Goodyera tesselata</i> Lodd.	.4	-		
Grasses	40.6	14.5		
<i>Maianthemum canadense</i> Desf.	.2	-		
<i>Melampyrum lineare</i> Desr.	.2	-		
<i>Potentilla tridentata</i> Soland.	.3	-		
<i>Streptopus roseus</i> Michx.	.2	-		
<i>Trifoliate borealis</i> Raf.	<.1	-		
Stratum total	49.9			
Moss				
<i>Calliergonella schreberi</i> (Willd., BR. & Sch.) Grout	7.1	5.2		
<i>Dicranum</i> Hedw. spp.	2.6	-		
<i>Polytrichum</i> Hedw. spp.	283.2	95.3		
All other mosses	360.4	118.5		
Stratum total	653.3			
Lichen				
<i>Cladonia mitis</i> Sandst.	939.1	325.7		
<i>Cladonia rangiferina</i> (L.) Web.	1,535.1	324.2		
<i>Peltigera aphthosa</i> (L.) Willd.	90.2	41.8		
All other lichens	743.8	161.6		
Stratum total	3,308.2			
Community total			4,137.0	

Table 6.--Aboveground oven-dry biomass by species and stratum for the aspen-birch-white pine community type of northeastern Minnesota

Tree Seedling	Little Sioux Fire						Cherokee-Frost Lake						Bucaw								
	n = 1			Foliage			n = 1			Foliage			n = 7			Foliage			n = 16		
	X	SE	Total	X	SE	Total	X	SE	Total	X	SE	Total	X	SE	Total	X	SE	Total	X	SE	Total
<i>Abies balsamea</i> (L.) Mill.																					
<i>Acer rubrum</i> L.																					
<i>Betula papyrifera</i> Marsh.																					
<i>Picea glauca</i> (Moench) Voss.																					
<i>Picea mariana</i> (Mill.) B.S.P.																					
<i>Pinus strobus</i> Lamb.																					
<i>Pinus taeda</i> L.																					
<i>Populus tremuloides</i> Michx.																					
<i>Populus grandidentata</i> Michx.																					
<i>Prunus pensylvanica</i> L.																					
<i>Stratum Total</i>																					
Sapling																					
<i>Abies balsamea</i> (L.) Mill.																					
<i>Acer rubrum</i> L.																					
<i>Betula papyrifera</i> Marsh.																					
<i>Picea mariana</i> (Mill.) B.S.P.																					
<i>Pinus strobus</i> Lamb.																					
<i>Pinus resinosa</i> Ait.																					
<i>Pinus strobus</i> L.																					
<i>Populus tremuloides</i> Michx.																					
<i>Sorbus americana</i> Marsh.																					
<i>Thuja occidentalis</i> L.																					
<i>Stratum Total</i>																					

Tree Seedling	Little Sioux Fire						Cherokee-Frost Lake						Bucaw								
	n = 1			Foliage			n = 1			Foliage			n = 7			Foliage			n = 16		
	X	SE	Total	X	SE	Total	X	SE	Total	X	SE	Total	X	SE	Total	X	SE	Total	X	SE	Total
<i>Abies balsamea</i> (L.) Mill.																					
<i>Acer rubrum</i> L.																					
<i>Betula papyrifera</i> Marsh.																					
<i>Fraxinus nigra</i> Marsh.																					
<i>Picea glauca</i> (Moench) Voss.																					
<i>Picea mariana</i> (Mill.) B.S.P.																					
<i>Pinus strobus</i> Lamb.																					
<i>Pinus resinosa</i> Ait.																					
<i>Pinus strobus</i> L.																					
<i>Populus tremuloides</i> Michx.																					
<i>Populus tremuloides</i> Michx.																					
<i>Sorbus americana</i> Marsh.																					
<i>Tilia cordata</i> L.																					
<i>Tall Shrub</i>																					
<i>Acer spicatum</i> Linn.																					
<i>Ailanthus altissima</i> (Ait.) Pursh.																					
<i>Cornus rugosa</i> Lam.																					
<i>Cornus stolonifera</i> Michx.																					
<i>Corylus cornuta</i> Marsh.																					
<i>Lonicera canadensis</i> Marsh.																					
<i>Lonicera tatarica</i> Linn.																					
<i>Stratum Total</i>																					

(Table 6 continued on next page)

(Table 6 continued on next page)

Table 6.—continued

	Little Sioux Fire												Cherokee-Frost Lake												
	n = 1						n = 1						n = 1						n = 7						n = 16
	Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		
	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	Total		
<i>Pyrola</i> L. spp.																									
<i>spicata</i> Nutt.																									
<i>hispidula</i> Michx.																									
<i>Streptopus</i> Zinn. spp.																									
<i>Tridentalis</i> Torrey & Raf.	1.4																								
<i>Trifolium</i> repens L.																									
<i>Trifolium pratense</i> L.																									
<i>Trifolium pratense</i> (Michx.) Salisb.																									
<i>Vicia americana</i> Michx.																									
<i>Viola</i> L. spp.																									
<i>Waldsteinia fragarioides</i> (Michx.) Tratt.	.1																								
Stratum total	534.0																								
Fern and fern allies																									
<i>Blechnum virginianum</i> (L.) Sm.																									
<i>Dryopteris spinulosa</i> (O. F. Muell.) Watt																									
<i>Lycopodium annotinum</i> L.																									
<i>Lycopodium clavatum</i> L.																									
<i>Lycopodium complanatum</i> L.	3.8																								
<i>Lycopodium lucidulum</i> Michx.																									
<i>Lycopodium obscurum</i> L.	12.7																								
<i>Osmunda claytoniana</i> L.																									
<i>Polyodium vulgare</i> L.																									
<i>Pteridium aquilinum</i> (L.) Kuhn.																									
<i>Roodsia villosa</i> (L.) R. Br.																									
Stratum total	16.5																								
Moss																									
<i>Calliergonella schreberi</i> (Willd.) Br. & Sch.-Burm.																									
Community total	17.05	T.3																							
	Little Sioux Fire												Cherokee-Frost Lake												
	n = 1						n = 1						n = 1						n = 7						n = 16
	Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		
	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	Total		
<i>Dicranum</i> Hedw. spp.																									
<i>obconicum</i> Spreng. (Hedw.) B. S. G.																									
<i>cristatum</i> & <i>castreum</i> (Hedw.)	0.2																								
<i>Polytrichum</i> Hedw. spp.																									
All other mosses																									
Stratum total	.2																								
Lichen																									
<i>Cladonia</i> alpestris (L.) Rabenh.																									
<i>Cladonia mitis</i> Sandst.																									
<i>Cladonia rangiferina</i> (L.) Neck.																									
<i>Cladonia sphagnicola</i> (L.) Hoffm.																									
<i>Peltigera aphitophora</i> (L.) Willd.																									
All other lichens																									
Stratum total	.2																								
Community total	1.05	T.3																							
	Little Sioux Fire												Cherokee-Frost Lake												
	n = 1						n = 1						n = 1						n = 7						n = 16
	Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		
	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	Total		
<i>Hypnum</i> Crat. & Castren. (Hedw.)																									
<i>lindbergii</i> (B. S. G.) B. S. G.																									
Stratum total	1.1																								
Lichen																									
<i>Cladonia</i> alpestris (L.) Rabenh.																									
<i>Cladonia mitis</i> Sandst.																									
<i>Cladonia rangiferina</i> (L.) Neck.																									
<i>Cladonia sphagnicola</i> (L.) Hoffm.																									
<i>Peltigera aphitophora</i> (L.) Willd.																									
All other lichens																									
Stratum total	.2																								
Community total	1.05	T.3																							
	Little Sioux Fire												Cherokee-Frost Lake												
	n = 1						n = 1						n = 1						n = 7						n = 16
	Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		
	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE	Total		
<i>Hypnum</i> Crat. & Castren. (Hedw.)																									
<i>lindbergii</i> (B. S. G.) B. S. G.																									
Stratum total	.2																								
Lichen																									
<i>Cladonia</i> alpestris (L.) Rabenh.																									
<i>Cladonia mitis</i> Sandst.																									
<i>Cladonia rangiferina</i> (L.) Neck.																									
<i>Cladonia sphagnicola</i> (L.) Hoffm.																									

Table 7.--Aboveground ovendry biomass by species and stratum for the aspen-birch community type of northeastern Minnesota

	(In Kg/ha)											
	Little Sioux Fire						Little Sioux Fire					
	Foliage		Total		Foliage		Total		Foliage		Total	
	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE
Tree												
<i>Abies balsamea</i> (L.) Mill.												
<i>Acer rubrum</i> L.												
<i>Betula papyrifera</i> Marsh.												
<i>Picea glauca</i> (Moench) Voss.												
<i>Picea mariana</i> (Mill.) B.S.P.												
<i>Pinus banksiana</i> Lamb.												
<i>Pinus resinosa</i> Ait.												
<i>Pinus strobus</i> L.												
<i>Populus grandidentata</i> Michx.												
<i>Populus tremuloides</i> Michx.												
<i>Thuja occidentalis</i> L.												
Stratum total												
Sapling												
<i>Abies balsamea</i> (L.) Mill.												
<i>Betula papyrifera</i> Marsh.												
<i>Fraxinus nigra</i> Marsh.												
<i>Picea glauca</i> (Moench) Voss.												
<i>Picea mariana</i> (Mill.) B.S.P.												
<i>Pinus banksiana</i> Lamb.												
<i>Pinus strobus</i> L.												
<i>Populus balsamifera</i> L.												
<i>Populus grandidentata</i> Michx.												
<i>Populus tremuloides</i> Michx.												
Stratum total												

	(In Kg/ha)											
	Little Sioux Fire						Little Sioux Fire					
	Foliage		Total		Foliage		Total		Foliage		Total	
	X	SE	X	SE	X	SE	X	SE	X	SE	X	SE
Tree seedling												
<i>Abies balsamea</i> (L.) Mill.	62.7	7.6	241.2	83.0	574.7	265.4	400.6	164.2	1,526.4	806.6		
<i>Acer rubrum</i> L.	2.4	-	3.2	-	7.9	-	207.0	-	785.4	-		
<i>Betula papyrifera</i> Marsh.												
<i>Fraxinus nigra</i> Marsh.												
<i>Picea glauca</i> (Moench) Voss.												
<i>Picea mariana</i> (Mill.) B.S.P.												
<i>Pinus banksiana</i> Lamb.	.2	.1	36.8	-	61.2	-	133.4	133.2	282.8	282.4		
<i>Pinus strobus</i> L.												
<i>Populus grandidentata</i> Michx.	110.4	-	165.4	3.1	704.2	95.8	671.8	324.1	3,094.7	1,141.9		
<i>Populus tremuloides</i> Michx.	843.6	692.2	543.8	519.4	1,075.2	966.3	1,112.4	570.1	7,360.2	4,773.6		
<i>Quercus rubra</i> L.												
<i>Sorbus americana</i> Marsh.												
Stratum total	1,019.3		990.4		2,423.2		2,525.2		13,049.5			
Tall shrub												
<i>Acer spicatum</i> Lam.	23.1	-										
<i>Alnus crispa</i> (Ait.) Pursh.												
<i>Alnus rugosa</i> (DuRoi) Spreng.												
<i>Amelanchier</i> Medic. spp.												
<i>Cornus rugosa</i> Lam.												
<i>Cornus stolonifera</i> Michx.												
<i>Corylus cornuta</i> Marsh.	45.2	34.6	391.4	74.7	1,253.2	494.2	347.0	120.1	1,266.5	590.1		
<i>Lonicera canadensis</i> Marsh.	19.1	-	20.7	20.2	44.4	43.4	69.6	64.6	169.1	160.1		
<i>Lonicera hirsuta</i> L.												
<i>Lonicera oblongifolia</i> (Goldie) Hook.												
<i>Lonicera villosa</i> (Michx.) R.D.S.												
<i>Prunus pensylvanica</i> L.F.	59.9	55.2	151.8	4.8	574.8	94.0	144.2	83.6	576.9	319.1		
<i>Prunus virginiana</i> L.												

Isabella Plantations										BWC AW					
6-10 n = 1				11-15 n = 1				21-25 n = 1				MATURE n = 10			
Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total	
\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE
48.6	1,397.5			71.3		955.4		109.3		1,537.2		362.6	153.0	2,802.9	1,182.0
2,288.6	21,140.7							356.7		8,928.7		607.6	169.1	15,449.0	4,166.5
								16.8		93.0		369.8	169.1	3,426.9	1,555.8
												300.4	195.9	2,478.9	1,523.7
371.5	6,758.6			203.1		1,402.8		1,235.8		21,379.2		511.9	363.2	13,509.8	9,986.0
	1,740.7					17,913.1		1,319.2		15,259.1		74.7	49.9	1,180.0	793.7
												186.2	158.3	4,288.9	3,679.5
.8	23.5											18.8	13.2	1,282.2	910.9
72.8	1,161.8			539.4		11,327.7		461.5		13,367.6		1,423.2	294.6	74,407.0	16,318.2
												4.9	-	38.8	-
2,782.3	30,482.1			2,554.5		31,599.0		3,499.3		60,564.8		3,860.3		II8,864.4	
												23.4	10.9	197.6	91.9
												114.9	20.2	4,200.9	746.5
												<.1	<.1	.3	.2
												22.3	10.5	147.3	68.8
												30.8	23.8	180.1	142.4
												1.8	1.0	14.1	8.0
												7.2	4.7	125.2	82.5
												.9	-	25.8	-
												1.6	.8	66.5	35.0
												145.4	37.2	3,750.4	1,008.2
												348.4		8,708.2	

(Table 7 continued on next page)

(Table 7.--continued)

	Little Sioux Fire											
	1 Foliage \bar{X} SE				3 Foliage \bar{X} SE				5 Foliage \bar{X} SE			
	n = 2	Total	n = 2	Total	n = 2	Total	n = 2	Total	n = 2	Total	n = 2	Total
<i>Salix bebbiana</i> Sarg.												
<i>Salix</i> L. spp.												
<i>Taxus canadensis</i> Marsh.												
<i>Viburnum rafinesquianum</i> Schult.												
Stratum Total		147.3		574.1		1,910.8		560.8		2,012.5		
Low Shrub												
<i>Arctostaphylos Uva-ursi</i> (L.) Spreng.												
<i>Chimaphila umbellata</i> (L.) Bart.												
<i>Comptonia peregrina</i> B. & B. Small. Rydb.	.1	0						37.0	-			
<i>Diervilla Lonicera</i> Mill.	.4	0						.2	-			
<i>Gaultheria procumbens</i> L.	.3	-										
<i>Ribes triste</i> Pall.												
<i>Ribes</i> L. spp.												
<i>Rosa blanda</i> Ait.												
<i>Rosa acicularis</i> Lindl.												
<i>Rubus parviflorus</i> Nutt.												
<i>Rubus pubescens</i> Raf.	.2	-						9.4	-			
<i>Rubus strigosus</i> Michx.	25.8	22.9						182.0	167.4			
<i>Vaccinium angustifolium</i> Ait.	7.2	-						25.4	23.8			
<i>Vaccinium myrtilloides</i> Michx.	.2	-						8.0	-			
Stratum Total		34.2						262.0				

	Little Sioux Fire											
	1 Foliage \bar{X} SE				3 Foliage \bar{X} SE				5 Foliage \bar{X} SE			
	n = 2	Total	n = 2	Total	n = 2	Total	n = 2	Total	n = 2	Total	n = 2	Total
Herb												
<i>Actaea rubra</i> (Ait.) Willd.												
<i>Anaphalis margaritacea</i> (L.) C.B. Clarke												
<i>Anemone quinquefolia</i> L.												
<i>Antennaria</i> spp. Gaertn.												
<i>Apocynum androsaemifolium</i> L.												
<i>Aralia hispida</i> Vent.	1.8	1.6						52.8	2.8			
<i>Aralia nudicaulis</i> L.	7.8	.8						37.4	2.1			
<i>Asarum canadense</i> L.												
<i>Aster ciliolatus</i> Lindl.												
<i>Aster macrophyllus</i> L.	235.3	53.8						139.4	29.6			
<i>Carex</i> L. spp.	13.1	1.2						71.4	31.1			
<i>Clintonia borealis</i> (Ait.) Raf.	4.4	2.0						55.2	32.4			
<i>Coptis groenlandica</i> (Oeder) Fern.	.1	-										
<i>Cornus canadensis</i> L.	30.0	3.6						67.7	11.6			
<i>Corydalis sempervirens</i> (L.) Pers.	9.5	1.6										
<i>Epilobium angustifolium</i> L.								21.7	-			
<i>Fragaria vesca</i> L.								.1	-			
<i>Fragaria virginiana</i> Duchesne.												
<i>Galium triflorum</i> Michx.	.2	-										
<i>Geranium bicknellii</i> Britt.	83.4	64.8						.1	-			
<i>Goodyera repens</i> (L.) R. Br.												
Grasses	.1	-						40.5	35.5			
<i>Hepatica americana</i> (D.C.) Ker.												
<i>Hieracium scabrum</i> Michx.												

Isabella Plantations												BWC AW			
6-10 n = 1				11-15 n = 1				21-25 n = 1				MATURE n = 10			
Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total	
\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE
0.1		0.2		100.2		1,127.6						1.2	0.8	5.4	4.0
346.6		3,063.4		396.4		3,234.1		503.3		888.8		403.8		22.0	14.3
10.3						21.0						.2	.1	.2	.1
36.2						53.1						.3		.4	
.7						27.6						3.6	2.2	22.0	14.3
16.4						93.9						107.4		34.4	
.1						140.7								<.1	
7.4						10.3								.2	
2.7						4.6								4.0	
73.8						351.2									

Isabella Plantations												BWC AW			
6-10 n = 1				11-15 n = 1				21-25 n = 1				MATURE n = 10			
Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total	
\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE
<0.1												0.2			
.5												.3	0.2		
												1.0	1.0		
												.6	.5		
												2.3	1.3		
.3												34.0	8.5		
												2.4	-		
												1.3	.9		
83.8				151.4				83.7				188.5	35.8		
<.1								25.6				8.3	2.9		
												9.7	3.2		
												4.0	3.0		
												7.3	2.3		
												2.4	1.6		
1.9												10.9	2.6		
3.0												.1	-		
												1.4	1.1		
105.7				149.9				68.9				.1	-		
												81.5	18.7		
												1.1	.9		
												.7	-		

(Table 7 continued on next page)

(Table 7.--continued)

	Little Sioux Fire											
	1 n = 2				3 n = 2				5 n = 2			
	Foliage		Total		Foliage		Total		Foliage		Total	
	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE
<u>Lactuca canadensis</u> L.												
<u>Lactuca</u> L. spp.												
<u>Lathyrus ochroleucus</u> Hook.	.1	-										
<u>Lathyrus venosus</u> Muhl.												
<u>Lathyrus</u> L. spp.												
<u>Linnaea borealis</u> L.	1.1	1.0			0.4	0.2			0.6	-		
<u>Lysimachia ciliata</u> L.												
<u>Maianthemum canadense</u> Desf.	1.6	.5			5.2	.4			13.6	4.6		
<u>Melampyrum lineare</u> Desr.												
<u>Mitella nuda</u> L.												
<u>Moneses uniflora</u> (L.) Gray												
<u>Petasites palmatus</u> (Ait.) Gray												
<u>Petasites vitifolius</u> Rydb.												
<u>Polygonatum pubescens</u> (Willd.) Pursh												
<u>Polygonum cilinode</u> Michx.	104.0	53.4			174.7	2.7			4.5	.3		
<u>Prunella vulgaris</u> L.												
<u>Pyrola elliptica</u> Nutt.												
<u>Pyrola rotundifolia</u> L.												
<u>Pyrola secunda</u> L.	<.1	-										
<u>Pyrola virens</u> Schweigg.												
<u>Pyrola</u> L. spp.												
<u>Sanicula marilandica</u> L.												
<u>Streptopus roseus</u> Michx.	.3	-			1.1	-			.9	-		
<u>Taraxacum</u> Zinn. spp.												
<u>Trientalis borealis</u> Raf.	.1	-			.6	.1			.1	-		
<u>Trifolium repens</u> L.												

	LITTLE SIOUX FIRE											
	1 n = 2				3 n = 2				5 n = 2			
	Foliage		Total		Foliage		Total		Foliage		Total	
	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE
<u>Uvularia sessilifolia</u> L.												
<u>Vicia americana</u> Muhl.												
<u>Viola</u> L. spp.	0.3	0.2			0.2	-			1.1	-		
Stratum total	493.3				668.5				327.1			
Fern and fern allies												
<u>Athyrium filix-femina</u> (L.) Roth.												
<u>Dryopteris spinulosa</u> (O.F. Muell.) Watt												
<u>Equisetum</u> L. spp.												
<u>Lycopodium annotinum</u> L.												
<u>Lycopodium clavatum</u> L.												
<u>Lycopodium complanatum</u> L.												
<u>Lycopodium obscurum</u> L.	3.2	3.1			12.2	-			19.4	-		
<u>Osmunda claytoniana</u> L.												
<u>Polypodium virginianum</u> L.												
<u>Pteridium aquilinum</u> (L.) Kuhn.												
Stratum total	3.2				12.2				20.6			
Moss												
<u>Calliergonella schreberi</u> (Willd., BR. & Sch.) Grout												
<u>Dicranum</u> Hedw. spp.												

	LITTLE SIOUX FIRE											
	1 n = 2				3 n = 2				5 n = 2			
	Foliage		Total		Foliage		Total		Foliage		Total	
	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE
<u>Polytrichum</u> Hedw. spp.	0.1	-			0.1	-			12.6	2.2		
All other mosses												
Stratum total	.1				.1				12.6			
Lichen												
<u>Cladonia mitis</u> Sandst.												
<u>Cladonia rangiferina</u> (L.) Web.												
All other lichens												
Stratum total												
Unidentified plants	.1	-										
Community total			1,697.5				5,276.8				15,821.1	

Isabella Plantations												BWC AW			
6-10 n = 1				11-15 n = 1				21-25 n = 1				MATURE n = 10			
Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total	
\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE
13.2		25.5		32.8				1.4				3.6	2.2		
5.0		2.3				1.6			5.3			1.5	1.4		
												4.5	2.6		
												.9	.7		
												.7	.3		
												2.1	1.6		
												.1	-		
												.1	.7		
												.1	-		
												1.0	-		
												.1	-		
												.2	.1		
												.2	-		
												.3	.2		
												<.1	-		
												.2	.1		
												1.8	-		
												1.4	.6		
												.2	.2		
												.5	.2		
												.1	-		

Isabella Plantations												BWC AW				
6-10 n = 1				11-15 n = 1				21-25 n = 1				MATURE n = 10				
Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		
\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	
213.5		368.4				219.6						385.5				
												2.0	1.9			
												.4	-			
												.2	.1			
												.2	-			
												.9	.6			
												.3	-			
												.2	-			
												2.7	2.0			
												3.3	-			
												1.9	-			
												93.7	19.3			
												105.3				
												43.8	19.9			
												2.9	1.6			

ISABELLA PLANTATIONS												BWC AW			
6-10 n = 1				11-15 n = 1				21-25 n = 1				MATURE n = 10			
Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total	
\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE
												21.8	13.1		
												298.3	52.7		
												366.8			
												4.8	-		
												11.6	8.4		
												27.5	11.1		
												43.9	-		
												1.2	-		
												34,293.3	37,215.1	65,567.6	131,829.6

Table 8.--Aboveground oven dry biomass by species and stratum for the fir-birch community type of northeastern Minnesota

Tree	Isabella Plantations (In Kg/ha)						Cherokee-Frost Lake n = 33						BWCAW MATURE n = 22									
	1-5 n = 2			11-15 n = 1			Foliage n = 7			Total n = 7			Foliage n = 7			Total n = 7			Foliage n = 22			
	\bar{X}	SE	Total	\bar{X}	SE	Total	\bar{X}	SE	Total	\bar{X}	SE	Total	\bar{X}	SE	Total	\bar{X}	SE	Total	\bar{X}	SE	Total	
<i>Abies balsamea</i> (L.) Mill.	2.3	-	21.5	-		130.0	1,061.6		256.9	139.5		2,168.6	1,194.5		2,385.3	427.0		18,805.5	3,362.4			
<i>Acer rubrum</i> L.																15.2	9.2		545.4	33.1		
<i>Betula papyrifera</i> Marsh.																1,479.9	333.1		34,537.8	7,731.6		
<i>Picea glauca</i> (Noeich) Voss.	45.1	-	350.9	-		1,117.2	27,228.5		1,107.9	157.5		31,922.8	6,888.2		1,004.4	253.6		9,640.4	2,685.5			
<i>Picea mariana</i> (Mill.) B.S.P.																1,287.3	397.2		11,637.8	3,546.5		
<i>Pinus strobus</i> L.	2.7	-	18.5	-		2,074.3	17,606.6		168.4	126.2		1,554.6	972.2		1,243.6	981.7		2,566.2	224.2			
<i>Pinus resinosa</i> Ait.	10.6	9.4	83.2	74.0		39,550.8	338.1		135.9	130.6		1,243.6	981.7		6,233.5			100.9				
<i>Pinus strobus</i> L.																			24.6	18.6		
<i>Populus tremuloides</i> Michx.	266.8	116.0	4,917.5	2,571.7		60.2	2,124.2		485.7	124.4		16,943.6	5,172.7		51.0	33.2		317.3	163.3			
<i>Sorbus americana</i> Marsh.																			6,649.0	3,443.6		
<i>Thuja occidentalis</i> L.																			181.8	34,18.8		
Stratum total	327.5		5,391.6			5,335.9			87,571.7	2,565.1		17.1	-		109.7	-		1,388.0	477.6			
Sapling																66,226.4	-		8,778.3	12,828.3		
<i>Abies balsamea</i> (L.) Mill.																						
<i>Acer rubrum</i> L.																						
<i>Betula papyrifera</i> Marsh.																						
<i>Picea glauca</i> (Noeich) Voss.																						
<i>Picea mariana</i> (Mill.) B.S.P.																						
<i>Pinus strobus</i> L.																						
<i>Populus tremuloides</i> Michx.																						
<i>Sorbus americana</i> Marsh.																						
<i>Thuja occidentalis</i> L.																						
Stratum total	1,149.1																25,829.2	1,467.7		13,503.6	1,467.7	

(Table 8 continued on next page)

(Table 8.--continued)

(Table 8 continued on next page)

(Table 8.--continued)

(Table 8 continued on next page)

(Table 8. --continued)

(Table 8 continued on next page)

(Table 8.--continued)

	Isabella Plantations						Cherokee-Frost Lake						BWCAW					
	1-5			11-15			33			MATURE			n = 22			Total		
	Foliage	n = 2	Total	Foliage	n = 1	Total	Foliage	n = 7	Total	Foliage	n = 22	Total	Foliage	n = 22	Total	Foliage	n = 22	Total
Fern and fern allies																		
<i>Athyrium filix-femina</i> (L.) Roth.																0.1	<.1	-
<i>Boltonia virginianum</i> (L.) Sw.																.2	0.2	
<i>Dryopteris spinulosa</i> (O.F. Muell.) Watt																.1	.1	
<i>Equisetum</i> L., spp.																3.0	-	
<i>Gymnocarpium dryopteris</i> (L.) Newm.																3.8	1.9	
<i>Lycopodium annotinum</i> L.																4.0	1.1	
<i>Lycopodium clavatum</i> L.																.2	.1	
<i>Lycopodium complanatum</i> L.																.1	.1	
<i>Lycopodium lucidulum</i> Michx.																.1	.1	
<i>Lycopodium obscurum</i> L.																4.0	2.0	
<i>Pteridium aquilinum</i> (L.) Kuhn.																2.6	1.1	
Stratum total		114.1	39.1					12.8								15.3		
Moss																		
<i>Calliergonella schreberi</i> (Wittd., Br. & Sch.) Grout																555.7	99.9	
<i>Dicranum</i> Hedw. spp.																43.1	11.4	
<i>Hypoleium sp. lendenii</i> (Hedw.) B.S.G.																16.9	8.7	
<i>Hypnum cristata-castrum</i> Hedw.																14.6	4.9	
<i>Polytrichum</i> Hedw. spp.																13.2	7.0	
<i>Sphagnum</i> L. spp.																6.4	-	
All other mosses																469.5	57.2	
Stratum total																395.4	119.4	
Lichen																		
<i>Cladonia rangiferina</i> (L.) Web.																3.4	3.0	
<i>Cladonia syringatica</i> (L.) Hoffm.																7.9	6.1	
All other lichens																14.2	4.9	
Stratum total																25.5		
Community total		6,397.4														91,469.5		
																95,972.5		
																	152,082.1	

Table 9.--Aboveground oven dry biomass by species and stratum for the maple-aspen-birch community type of northeastern Minnesota

Stratum/Species	(In Kg/ha)			
	BWC AW		MATURE (n = 20)	
	Foliage	Total		
Tree				
<i>Abies balsamea</i> (L.) Mill.	278.8	89.6	2,165.4	694.1
<i>Acer rubrum</i> L.	471.5	90.0	18,646.1	3,714.4
<i>Betula papyrifera</i> Marsh.	906.0	160.2	22,950.0	3,986.2
<i>Fraxinus nigra</i> Marsh.	.2	-	53.0	-
<i>Picea glauca</i> (Moench) Voss.	200.2	105.2	1,833.1	972.2
<i>Picea mariana</i> (Mill.) B.S.P.	103.1	36.7	948.2	344.5
<i>Pinus banksiana</i> Lamb.	179.7	75.1	4,336.7	1,717.4
<i>Pinus resinosa</i> Ait.	890.7	449.1	12,677.8	6,378.2
<i>Pinus strobus</i> L.	949.2	511.1	21,823.7	11,554.8
<i>Populus grandidentata</i> Michx.	94.9	38.9	6,610.4	2,811.7
<i>Populus tremuloides</i> Michx.	1,057.7	151.0	56,737.4	8,429.8
<i>Quercus rubra</i> L.	77.6	44.2	1,560.8	921.2
<i>Sorbus americana</i> Marsh.	2.3	-	26.4	-
Stratum total	5,214.0		15,0369.0	
Sapling				
<i>Abies balsamea</i> (L.) Mill.	136.2	63.3	1,151.5	533.6
<i>Acer rubrum</i> L.	247.4	58.1	4,568.0	1,097.3
<i>Betula papyrifera</i> Marsh.	152.6	25.6	5,495.4	908.3
<i>Fraxinus nigra</i> Marsh.	.9	.7	11.4	8.1
<i>Picea glauca</i> (Moench) Voss.	21.6	13.6	137.8	87.0
<i>Picea mariana</i> (Mill.) B.S.P.	22.3	15.3	136.5	99.2
<i>Pinus resinosa</i> Ait.	1.2	.8	11.8	7.5
<i>Pinus strobus</i> L.	.6	.5	11.6	8.4
<i>Populus grandidentata</i> Michx.	5.7	2.1	235.8	84.4
<i>Populus tremuloides</i> Michx.	57.8	15.0	1,499.7	390.2
<i>Quercus rubra</i> L.	9.7	5.8	96.7	57.9
<i>Sorbus americana</i> Marsh.	5.4	2.6	56.3	27.4
Stratum total	661.4		13,412.5	
Tree seedling				
<i>Abies balsamea</i> (L.) Mill.	30.6	12.2	69.5	28.5
<i>Acer rubrum</i> L.	12.0	2.0	169.4	30.2
<i>Betula papyrifera</i> Marsh.	.9	.2	20.4	5.3
<i>Fraxinus nigra</i> Marsh.	1.0	.6	5.5	3.6
<i>Picea glauca</i> (Moench) Voss.	5.4	2.9	10.1	5.7
<i>Picea mariana</i> (Mill.) B.S.P.	.6	.4	1.0	.6
<i>Pinus resinosa</i> Ait.	.1	.1	.5	.3
<i>Pinus strobus</i> L.	2.1	.8	13.1	4.5
<i>Populus grandidentata</i> Michx.	.2	.2	1.3	1.0
<i>Populus tremuloides</i> Michx.	8.7	1.6	51.3	10.9
<i>Quercus rubra</i> L.	7.9	3.4	37.7	17.2
<i>Sorbus americana</i> Marsh.	.8	.4	4.0	1.8
<i>Thuja occidentalis</i> L.	.2	-	.4	-
Stratum total	70.5		384.2	
Tall shrub				
<i>Acer spicatum</i> Lam.	49.9	15.6	397.9	139.5
<i>Alnus crispa</i> (Ait.) Pursh.	38.0	19.4	286.2	147.3
<i>Amelanchier</i> Medic. spp.	11.4	4.4	112.7	41.5
<i>Cornus rugosa</i> Lam.	9.3	5.8	47.3	36.3
<i>Lonicera canadensis</i> Marsh.	1.1	.5	5.6	3.4
<i>Lonicera villosa</i> (Michx.) R.D.S.	<.1	-	<.1	-
<i>Prunus pensylvanica</i> L.F.	<.1	-	<.1	-
<i>Prunus virginiana</i> L.	.1	.1	.5	.5
<i>Salix bebbiana</i> Sarg.	13.7	13.2	532.1	530.0
<i>Salix</i> L. spp.	<.1	-	<.1	-
<i>Viburnum regnesquianum</i> Schult.	.8	.6	2.8	2.1
Stratum total	331.0		2,644.7	

(Table 9 continued on next page)

(Table 9.--continued)

Stratum/Species	BWC AW MATURE (n = 20)		Total	
	Foliage		X	SE
Low shrub				
<u>Arctostaphylos Uva-ursi</u> (L.) Spreng.		0.4		-
<u>Chimaphila umbellata</u> (L.) Bart.		.2		0.1
<u>Comptonia peregrina</u> B.&B. Small. Rydb.		1.4		1.0
<u>Diervilla Lonicera</u> Mill.		11.4		2.2
<u>Gaultheria procumbens</u> L.		.8		.4
<u>Rosa acicularis</u> Lindl.		.1		-
<u>Rubus pubescens</u> Raf.		1.4		.8
<u>Rubus strigosus</u> Michx.		1.2		.4
<u>Vaccinium angustifolium</u> Ait.		9.0		2.9
<u>Vaccinium myrtilloides</u> Michx.		8.3		5.9
Stratum total		34.2		
Herb				
<u>Anaphalis margaritacea</u> (L.) C.B. Clarke	0.1			
<u>Anemone quinquefolia</u> L.	.1	0.1		
<u>Apocynum androsaemifolium</u> L.	1.0	.8		
<u>Aralia nudicaulis</u> L.	27.4	4.6		
<u>Aster ciliolatus</u> Lindl.	.2			
<u>Aster macrophyllum</u> L.	132.7	20.0		
<u>Carex</u> L. spp.	10.7	5.5		
<u>Clintonia borealis</u> (Ait.) Raf.	11.6	2.9		
<u>Coptis groenlandica</u> (Oeder) Fern.	1.7	.8		
<u>Cornus canadensis</u> L.	8.9	3.6		
<u>Epilobium angustifolium</u> L.	.2			
<u>Fragaria vesca</u> L.	2.3	1.2		
<u>Galium triflorum</u> Michx.	.1	<.1		
<u>Geranium bicknellii</u> Britt.	.2	.1		
<u>Goodyera repens</u> (L.) R. Br.	.2	.1		
<u>Goodyera tesselata</u> Lodd.	.1	.1		
Grasses	26.9	5.3		
<u>Impatiens capensis</u> Meerb.	.3	.2		
<u>Lathyrus ochroleucus</u> Hook.	.7	.3		
<u>Lathyrus venosus</u> Muhl.	.1			
<u>Linnaea borealis</u> L.	4.1	1.6		
<u>Maianthemum canadense</u> Desf.	8.6	1.2		
<u>Melampyrum lineare</u> Desr.	.5	.3		
<u>Mitella nuda</u> L.	.1			
<u>Monotropa uniflora</u> L.	<.1			
<u>Petasites vitifolius</u> Rydb.	.3			
<u>Pyrola rotundifolia</u> L.	<.1			
<u>Pyrola secunda</u> L.	.1	.1		
<u>Streptopus roseus</u> Michx.	2.0	.5		
<u>Trientalis borealis</u> Raf.	.6	.1		
<u>Uvularia sessilifolia</u> L.	.2	.1		
<u>Viola</u> L. spp.	.5	.1		
Stratum total	242.6			
Fern and fern allies				
<u>Dryopteris Phegopteris</u> L.	.5			
<u>Dryopteris spinulosa</u> (O.F. Muell.) Watt	1.2	1.1		
<u>Gymnocarpium Dryopteris</u> (L.) Newm.	.3			
<u>Lycopodium annotinum</u> L.	3.1	1.4		
<u>Lycopodium clavatum</u> L.	4.2	1.3		
<u>Lycopodium complanatum</u> L.	1.9	1.9		
<u>Lycopodium obscurum</u> L.	9.7	2.2		
<u>Osmunda claytoniana</u> L.	2.0	1.8		
<u>Polyodium virginianum</u> L.	.3			
<u>Pteridium aquilinum</u> (L.) Kuhn.	51.7	13.4		
Stratum total	74.9			
Moss				
<u>Calliergonella schreberi</u> (Willd., BR. & Sch.) Grout	84.3	21.8		
<u>Dicranum Hedw.</u> spp.	27.0	14.1		
<u>Hypnum crista-castrrensis</u> Hedw.	.7			
<u>Polytrichum Hedw.</u> spp.	11.4	8.1		
All other mosses	304.4	48.4		
Stratum total	427.8			
Lichen				
<u>Cladonia mitis</u> Sandst.	.5			
<u>Cladonia rangiferina</u> (L.) Web.	7.4			
<u>Cladonia sylvatica</u> (L.) Hoffm.	.5			
<u>Peltigera aphthosa</u> (L.) Willd.	1.0	.7		
All other Lichens	25.7	6.8		
Stratum total	35.1			
Community total	167,625.0			

Table 10.--Aboveground oven dry biomass by species and stratum for the maple-aspen-birch-fir community type of northeastern Minnesota

Stratum/Species Tree	(In Kg/ha)			
	BWC AW			
	MATURE (n = 19)			
	Foliage			Total
X	SE	X	SE	
<i>Abies balsamea</i> (L.) Mill.	1,402.8	376.3	11,054.6	2,987.3
<i>Acer rubrum</i> L.	425.8	103.5	16,099.1	3,957.9
<i>Betula papyrifera</i> Marsh.	1,084.3	303.8	24,927.6	7,103.9
<i>Picea glauca</i> (Moench) Voss.	289.8	75.7	2,637.4	714.8
<i>Picea mariana</i> (Mill.) B.S.P.	594.3	176.4	5,465.3	1,696.8
<i>Pinus banksiana</i> Lamb.	174.6	104.6	4,530.3	2,922.9
<i>Pinus resinosa</i> Ait.	320.6	224.6	5,051.0	3,515.2
<i>Pinus strobus</i> L.	982.7	423.6	23,497.5	10,136.9
<i>Populus grandidentata</i> Michx.	.34.3	20.1	2,395.0	1,382.9
<i>Populus tremuloides</i> Michx.	1,296.6	301.2	73,057.3	18,290.4
<i>Thuja occidentalis</i> L.	.90.4	88.8	1,408.9	1,392.4
Stratum total	6,696.2		170,124.0	
Sapling				
<i>Abies balsamea</i> (L.) Mill.	567.8	149.3	4,783.8	1,255.3
<i>Acer rubrum</i> L.	206.1	44.0	3,966.0	832.0
<i>Betula papyrifera</i> Marsh.	.75.7	30.4	2,728.5	1,085.7
<i>Picea glauca</i> (Moench) Voss.	41.2	22.8	272.2	153.9
<i>Picea mariana</i> (Mill.) B.S.P.	129.5	53.3	756.2	296.7
<i>Pinus banksiana</i> Lamb.	.6	-	4.8	-
<i>Pinus resinosa</i> Ait.	.6	.4	6.1	4.2
<i>Pinus strobus</i> L.	.5	.4	9.4	7.8
<i>Populus grandidentata</i> Michx.	.4	.4	15.8	15.3
<i>Populus tremuloides</i> Michx.	37.2	12.7	1,014.5	352.2
<i>Quercus rubra</i> L.	.9	-	5.8	-
<i>Sorbus americana</i> Marsh.	24.9	18.8	246.5	184.2
Stratum total	1,085.4		13,809.6	
Tree seedling				
<i>Abies balsamea</i> (L.) Mill.	50.3	6.9	114.1	16.2
<i>Acer rubrum</i> L.	13.2	2.9	203.2	55.0
<i>Betula papyrifera</i> Marsh.	1.3	.2	27.6	5.6
<i>Fraxinus nigra</i> Marsh.	.1	-	.5	-
<i>Picea glauca</i> (Moench) Voss.	3.3	1.4	5.6	2.3
<i>Picea mariana</i> (Mill.) B.S.P.	4.6	2.6	7.9	4.4
<i>Pinus resinosa</i> Ait.	<.1	-	.2	-
<i>Pinus strobus</i> L.	1.5	.6	10.2	4.0
<i>Populus grandidentata</i> Michx.	.3	-	1.6	-
<i>Populus tremuloides</i> Michx.	10.2	2.1	57.0	12.4
<i>Quercus rubra</i> L.	3.4	3.1	18.5	16.8
<i>Sorbus americana</i> Marsh.	6.2	1.7	33.9	9.5
<i>Thuja occidentalis</i> L.	.1	-	.2	-
<i>Tilia americana</i> L.	<.1	-	.2	-
Stratum total	94.6		480.7	
Tall shrub				
<i>Acer spicatum</i> Lam.	96.8	29.5	867.6	313.3
<i>Alnus crispa</i> (Ait.) Pursh.	.6	.4	4.8	4.4
<i>Alnus rugosa</i> (DuRoi) Spreng.	<.1	-	<.1	-
<i>Amelanchier</i> Medic. spp.	16.4	14.9	127.2	110.2
<i>Cornus rugosa</i> Lam.	.6	.4	2.8	1.9
<i>Corylus cornuta</i> Marsh.	95.0	25.3	565.6	162.8
<i>Lonicera canadensis</i> Marsh.	.9	.3	3.5	1.3
<i>Lonicera villosa</i> (Michx.) R.d.S.	<.1	<.1	<.1	<.1
<i>Prunus pensylvanica</i> L.F.	13.4	11.7	75.9	65.1
<i>Prunus virginiana</i> L.	.1	.1	1.3	1.3
<i>Salix bebbiana</i> Sarg.	<.1	-	<.1	-
<i>Salix</i> L. spp.	<.1	-	<.1	-
<i>Viburnum rafinesquianum</i> Schult.	1.9	1.9	18.9	18.8
Stratum total	225.8		1,667.8	

(Table 10 continued on next page)

(Table 10.--continued)

Stratum/species	BWCW MATURE (n = 19)		Total	
	<u>Foliage</u>	<u>X</u>	<u>X</u>	<u>SE</u>
Low shrub				
<i>Chimaphila umbellata</i> (L.) Bart.		0.6	0.3	
<i>Dierivilla Lonicera</i> Mill.		6.4	2.1	
<i>Gaultheria hispidula</i> (L.) Muhl.		.7	.5	
<i>Gaultheria procumbens</i> L.		3.5	1.8	
<i>Ledum groenlandicum</i> Oeder.		<.1	-	
<i>Ribes</i> L. spp.		.1	-	
<i>Rosa acicularis</i> Lindl.		.8	.7	
<i>Rubus pubescens</i> Raf.		3.3	1.8	
<i>Rubus strigosus</i> Michx.		11.1	7.1	
<i>Vaccinium angustifolium</i> Ait.		10.6	2.8	
<i>Vaccinium myrtilloides</i> Michx.		4.9	2.2	
Stratum total		42.1		
Herb				
<i>Anemone quinquefolia</i> L.	0.2	0.1		
<i>Aralia nudicaulis</i> L.	22.6	4.7		
<i>Aster ciliolatus</i> Lindl.	.1	-		
<i>Aster macrophyllus</i> L.	83.2	13.0		
<i>Carex</i> L. spp.	5.2	2.0		
<i>Circaeal alpina</i> L.	.2	-		
<i>Clintonia borealis</i> (Ait.) Raf.	17.0	3.6		
<i>Coptis groenlandica</i> (Oeder) Fern.	2.6	.8		
<i>Cornus canadensis</i> L.	20.8	3.6		
<i>Epilobium angustifolium</i> L.	.1	-		
<i>Epigaea repens</i> L.	.1	-		
<i>Fragaria vesca</i> L.	1.1	.7		
<i>Fragaria virginiana</i> Duchesne.	.2	.2		
<i>Galium triflorum</i> Michx.	.5	.2		
<i>Goodyera repens</i> (L.) R. Br.	.1	.1		
<i>Goodyera tesselata</i> Lodd.	.1	.1		
Grasses	19.4	3.2		
<i>Habenaria orbiculata</i> (Pursh) Torr.	.2	-		
<i>Habenaria</i> spp.	<.1	-		
<i>Lathyrus ochroleucus</i> Hook.	.3	-		
<i>Linnaea borealis</i> L.	4.0	1.1		
<i>Maianthemum canadense</i> Desf.	12.4	1.2		
<i>Metamprum lineare</i> Desr.	.6	.2		
<i>Mitella nuda</i> L.	.3	-		
<i>Petasites vitifolius</i> Rydb.	.4	.3		
<i>Polygonum ciliinode</i> Michx.	.7	.4		
<i>Pyrola secunda</i> L.	.2	.1		
<i>Pyrola virens</i> Schweigg.	<.1	-		
<i>Solidago</i> L. spp.	.2	.1		
<i>Streptopus roseus</i> Michx.	2.2	.7		
<i>Trientalis borealis</i> Raf.	.8	.1		
<i>Trifolium repens</i> L.	<.1	-		
<i>Viola</i> L. spp.	.9	.2		
<i>Waldsteinia fragarioides</i> (Michx.) Tratt.	.1			
Stratum total	196.9			
Fern and fern allies				
<i>Athyrium filix-femina</i> (L.) Roth.	1.8	-		
<i>Dryopteris spinulosa</i> (O.F. Muell.) Watt	.4	.3		
<i>Equisetum</i> L. spp.	<.1	-		
<i>Gymnocarpium Dryopteris</i> (L.) Newm.	.2	.1		
<i>Lycopodium annotinum</i> L.	5.0	1.6		
<i>Lycopodium clavatum</i> L.	11.8	5.6		
<i>Lycopodium complanatum</i> L.	.6	.4		
<i>Lycopodium Tucidulum</i> Michx.	2.9	-		
<i>Lycopodium obscurum</i> L.	12.4	2.6		
<i>Osmunda claytoniana</i> L.	1.9	-		
<i>Polypodium virginianum</i> L.	.3	.2		
<i>Pteridium aquilinum</i> (L.) Kuhn.	34.3	9.5		
Stratum total	71.7			
Moss				
<i>Calliergonella schreberi</i>	207.6	50.8		
(Willd., BR. & Sch.) Grout				
<i>Dicranum</i> Hedw. spp.	36.6	13.2		
<i>Hylocomium splendens</i> (Hedw.) B.S.G.	.2	-		
<i>Hypnum cristata-castrensis</i> Hedw.	2.2	2.0		
<i>Polytrichum</i> Hedw. spp.	9.3	2.7		
<i>Sphagnum</i> L. spp.	2.0	-		
All other mosses	287.0	42.9		
Stratum total	544.9			
Lichen				
<i>Cladonia rangiferina</i> (L.) Web.	10.0	4.6		
<i>Cladonia sylvatica</i> (L.) Hoffm.	7.5	5.8		
<i>Peltigera aphthosa</i> (L.) Willd.	.7	-		
All other Lichens	17.0	7.1		
Stratum total	35.2			
Community total	186,972.9			

Table 11.--Aboveground ovendry biomass by species and stratum for the jack pine-oak community type of northeastern Minnesota

Tree	(In Kg/ha)											
	Little Sioux Fire						n = 1					
	Foliage		Total		Foliage		Total		Foliage		Total	
	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE
Abies balsamea (L.) Mill.												
Acer rubrum L.												
Betula papyrifera Marsh.												
Picea glauca (Moench) Voss.												
Picea mariana (Mill.) B.S.P.												
Pinus banksiana Lamb.												
Pinus resinosa Ait.												
Pinus strobus L.												
Populus grandidentata Michx.												
Populus tremuloides Michx.												
Quercus rubra L.												
Stratum total												
Sapling												
Abies balsamea (L.) Mill.												
Acer rubrum L.												
Betula papyrifera Marsh.												
Picea glauca (Moench) Voss.												
Picea mariana (Mill.) B.S.P.												
Pinus banksiana Lamb.												
Pinus resinosa Ait.												
Pinus strobus L.												
Populus grandidentata Michx.												
Populus tremuloides Michx.												
Quercus rubra L.												
Stratum total												

Tree seedling	(In Kg/ha)											
	Little Sioux Fire						n = 1					
	Foliage		Total		Foliage		Total		Foliage		Total	
	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE
Abies balsamea (L.) Mill.												
Acer rubrum L.					21.4		61.7		130.4		269.0	
Betula papyrifera Marsh.											421.0	
Picea mariana (Mill.) B.S.P.												1,883.8
Pinus banksiana Lamb.					1.4		62.2		91.0		904.3	
Pinus strobus L.												1,737.8
Populus balsamifera L.												
Populus grandidentata Michx.												
Populus tremuloides Michx.												
Quercus rubra L.												
Sorbus americana Marsh.												
Stratum total					24.3		213.1		483.1		1,671.9	
Tall shrub												
Alnus crispa (Ait.) Pursh.												
Amelanchier Medic. spp.												
Cornus rugosa Lam.					13.7		20.6		43.6		302.8	
Corylus cornuta Marsh.												
Lonicera canadensis Marsh.												
Lonicera oblongifolia (Goldie) Hook.												
Lonicera villosa (Michx.) R.dS.												
Prunus pensylvanica L.F.												
Prunus virginiana L.												
Salix bebbiana Sarg.												
Salix L. spp.												
Spiraea L. spp.												
Stratum total					90.2						147.3	
												605.0
					188.5		52.7		121.6		968.5	
												3,414.7

Isabella Plantations												BWC AW				
11-15 n = 2				21-25 n = 2				26-30 n = 1				MATURE n = 6				
Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		
\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	
50.2	-	384.9	-	1,281.0	-	9,900.6	-			10.8	-	87.3	-			
102.4	50.6	2,497.4	861.7	93.8	17.9	2,564.2	192.5			9.1	5.9	252.4	163.2			
390.6	177.1	3,042.4	1,337.4	838.5	817.9	7,218.2	7,086.4			22.3	15.2	596.8	399.4			
459.8	-	3,942.2	-	548.9	94.6	4,357.7	1,443.3	47.7	237.6	583.4	447.4	5,015.9	3,888.0			
2,432.1	1,601.1	24,572.3	15,871.4	2,151.3	-	30,665.5	-	2,645.7	37,501.9	5,230.2	783.3	95,630.2	13,837.6			
507.8	-	4,886.9	-	2,256.0	763.8	24,902.4	8,571.5	3,556.6	40,409.6	423.0	399.0	6,363.2	6,021.9			
				177.5	-	3,677.2	-			286.5	-	6,263.0	-			
					3.0	.6	73.9	29.1	577.5	25,657.7	18.8	15.8	1,283.7	1,114.1		
										30.1	11.1	1,292.2	559.1			
										2.4	-	41.6	-			
3,942.9		39,326.1		7,350.0		83,359.7		6,827.5		103,806.8		6,616.6		116,826.3		
												3.0	2.4	26.0	20.0	
												18.0	15.7	353.0	319.0	
												16.8	6.6	644.9	247.0	
												12.7	12.2	83.1	80.6	
												215.2	108.8	1,215.1	623.3	
												573.5	341.7	5,730.6	3,326.6	
												1.0	-	8.4	-	
												2.4	1.3	41.9	22.3	
												.3	-	11.3	-	
												.8	.7	20.5	18.3	
												10.9	4.6	98.4	42.0	
												854.6			8,233.2	

Isabella Plantations												BWC AW			
11-15 n = 2				21-25 n = 2				26-30 n = 1				MATURE n = 6			
Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total	
\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE
				<0.1	-	<0.1	-			3.2	2.1	7.3	4.8		
				2.6	2.5	57.8	57.6			1.9	.5	35.0	9.2		
										.6	.4	15.5	10.6		
										36.6	22.6	68.3	42.9		
										.1	-	.6	-		
										14.4	13.8	126.3	121.7		
				<.1	-	.6	-					2.4	1.4	16.0	9.4
												7.2	3.6	38.6	19.8
												.1	.7	-	
												66.5	-	308.3	
				2.7		58.5									
77.0	-	669.8	-	74.9	-	779.9	-	194.8	-	1,905.7	-	1.0	-	4.0	-
13.2	11.1	86.6	76.8	17.2	4.8	149.5	64.5	7.3	47.6		.6	.4	12.0	5.7	
.2	-	.6	-					16.2	81.4						
76.0	69.1	693.2	631.2	124.6	-	1,233.7	-	178.1	-	1,494.9	-	18.2	7.0	95.7	43.4
4.0	3.0	30.6	22.3	3.3	2.7	22.1	17.5								
.3	-	1.1	-	<.1	-	<.1	-	2.6		10.6					
.6	-	4.0	-												
.8	.4	9.2	4.6	8.0	3.3	78.6	29.8								
				7.6	-	52.0	-								
				.5	-	1.6	-	8.6	-	37.5	-	5.1	-	40.5	-
												.3	-	.6	-
172.1		1,495.1		236.2		2,317.5		407.6		3,577.7		30.1		226.7	

(Table 11.--continued)

	Little Sioux Fire											
	1 n = 1				3 n = 1				5 n = 1			
	Foliage		Total		Foliage		Total		Foliage		Total	
	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE
Low shrub												
<i>Chimaphila umbellata</i> (L.) Bart.												
<i>Comptonia peregrina</i> B.&B. Small. Rydb.	40.5								799.4			1,377.0
<i>Diervilla lonicera</i> Mill.	4.8								3.6			28.4
<i>Gaultheria hispida</i> (L.) Muhl.												
<i>Gaultheria procumbens</i> L.	8.6								52.2			44.5
<i>Ledum groenlandicum</i> Oeder.												
<i>Ribes</i> L. spp.												
<i>Rosa blanda</i> Ait.												
<i>Rosa acicularis</i> Lindl.												
<i>Rubus pubescens</i> Raf.												
<i>Rubus strigosus</i> Michx.												
<i>Vaccinium angustifolium</i> Ait.	46.2								290.0			54.5
<i>Vaccinium myrtilloides</i> Michx.	15.0								8.7			156.6
Stratum total	115.1								1,153.9			1,661.0
Herb												
<i>Anemone quinquefolia</i> L.												
<i>Apocynum androsaemifolium</i> L.												
<i>Aralia hispida</i> Vent.	1.1								203.3			103.7
<i>Aralia nudicaulis</i> L.	12.7								49.8			24.1
<i>Aster macrophyllus</i> L.	333.3								194.1			135.0
<i>Carex</i> L. spp.	14.4								24.7			
<i>Clintonia borealis</i> (Ait.) Raf.	1.6											
<i>Coptis groenlandica</i> (Oeder) Fern.												
<i>Cornus canadensis</i> L.	.1								6.8			
<i>Corydalis sempervirens</i> (L.) Pers.	1.9											
	Little Sioux Fire											
	1 n = 1				3 n = 1				5 n = 1			
	Foliage		Total		Foliage		Total		Foliage		Total	
	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE
Herb												
<i>Epilobium angustifolium</i> L.												7.7
<i>Epigaea repens</i> L.												
<i>Fragaria vesca</i> L.												
<i>Fragaria virginiana</i> Duchesne.												
<i>Geranium bicknellii</i>	0.3											
<i>Goodyera repens</i> (L.) R. Br.												
<i>Goodyera tesselata</i> Lodd.												
Grasses									251.7			46.0
<i>Hieracium</i> L. spp.												
<i>Lathyrus ochroleucus</i> Hook.												
<i>Linnaea borealis</i> L.												
<i>Maianthemum canadense</i> Desf.	3.9								14.0			13.6
<i>Melampyrum lineare</i> Desr.									1.6			
<i>Polygonia paucifolia</i> Willd.												
<i>Polygonum cilinode</i> Michx.	2.5								14.4			.7
<i>Pyrola rotundifolia</i> L.												
<i>Pyrola secunda</i> L.												
<i>Pyrola virens</i> Schweigg.												
<i>Solidago</i> L. spp.												
<i>Streptopus roseus</i> Michx.												
<i>Trientalis borealis</i> Raf.												
<i>Viola</i> L. spp.	<.1											
Stratum total	371.9								760.4			330.8
Fern and fern allies												
<i>Lycopodium clavatum</i> L.												
<i>Lycopodium complanatum</i> L.												
<i>Lycopodium obscurum</i> L.	.8											
<i>Pteridium aquilinum</i> (L.) Kuhn.	301.7								961.5			264.8
Stratum total	302.5								961.5			264.8
	Little Sioux Fire											
	1 n = 1				3 n = 1				5 n = 1			
	Foliage		Total		Foliage		Total		Foliage		Total	
	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE
Moss												
<i>Calliergonella schreberi</i>												
(Willd., Br. & Sch.) Grout												
<i>Dicranum</i> Hedw. spp.												
<i>Polytrichum</i> Hedw. spp.	0.1								5.2			33.2
All other mosses												
Stratum total	.1								5.2			33.2
Lichen												
<i>Cladonia mitis</i> Sandst.												
<i>Cladonia rangiferina</i> (L.) Web.												
All other lichens												
Stratum total												
Unidentified plants												
Community total												
	1,002.5				3,485.8				10,423.3			

Isabella Plantations										BWC AW					
11-15 n = 2				21-25 n = 2				26-30 n = 1				MATURE n = 6			
Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total	
\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE
2.9	-					1.2	-							1.8	-
61.1	-									21.6				2.7	1.8
68.9	61.1					20.4	17.1			103.2				33.3	14.0
						3.7	-							3.1	1.6
69.2	33.3					66.0	32.5			29.8				67.6	10.9
						17.6	-								
										1.1					
9.2	-														
60.5	43.4					20.0	6.6			29.4				1.8	1.2
.3	.3					.5	.5			.8					
19.4	1.6					4.1	.9			20.0					
82.0	46.1					12.3	1.1			28.1				198.4	33.6
89.1	32.3					74.9	57.0							14.5	8.4
462.6						220.7				234.0				323.2	

7.5	-	<0.1	-			<0.1						<0.1	1.0	0.7
1.8	1.7	.7	-			7.0						5.7	2.0	
76.0	38.0	17.9	14.1			106.3						68.5	26.1	

Isabella Plantations										BWC AW					
11-15 n = 2				21-25 n = 2				26-30 n = 1				MATURE n = 6			
Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total	
\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE
5.4	-	1.3	-	.4	-			9.8				0.7	.3	-	
		.4	-												
133.2	54.1	77.4	32.6			57.3						44.2	12.8		
2.5	-					<.1									
8.6	-	7.0	2.0			25.4						17.5	10.0		
8.4	1.5	1.5	1.2			9.1						12.9	2.1		
<.1	-											5.4	3.5		
<.1	0	<.1	-			<.1									
2.7	-	1.1	-			3.4						.1	.1		
2.7	-	.1	-			5.7									
1.1	-	1.1	-			<.1									
		<.1	-			<.1									
		<.1	-												
296.4		.4	-									177.6			
		137.2				225.9									
<.1	-					2.7						.2	-		
		<.1	-			<.1									
<.1	-											.3	.2		
57.9	41.2	16.7	0			117.7						90.0	48.9		
58.0		16.8				120.5						90.5	18.4		

Isabella Plantations										BWC AW					
11-15 n = 2				21-25 n = 2				26-30 n = 1				MATURE n = 6			
Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total	
\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE
												672.1	232.3		
												254.9	47.8		
												74.2	49.5		
												137.2	65.5		
												1,138.4			
												282.2	209.7		
												247.8	100.6		
												116.2	70.7		
												646.2	78.1		

41,638.2

86,110.4

107,964.9

127,970.4

Table 12.--Aboveground oven dry biomass by species and stratum for the maple-oak community type of northeastern Minnesota

Stratum/species	(In Kg/ha)			
	BWC AW		MATURE	
			(n = 4)	
	Foliage		Total	
Tree				
Abies balsamea (L.) Mill.	1,087.0	986.8	8,558.0	7,755.5
Acer rubrum L.	515.8	304.9	17,127.7	10,869.8
Betula papyrifera Marsh.	141.7	90.9	3,798.5	2,455.5
Picea glauca (Moench) Voss.	172.6	-	1,444.0	-
Picea mariana (Mill.) B.S.P.	66.9	-	546.3	-
Pinus banksiana Lamb.	89.8	-	1,944.1	-
Pinus resinosa Ait.	1,839.8	1,170.2	26,246.1	16,966.7
Pinus strobus L.	327.0	211.5	6,993.6	4,472.1
Populus grandidentata Michx.	89.6	65.9	5,396.8	3,832.4
Populus tremuloides Michx.	547.8	236.9	25,291.7	11,520.3
Quercus rubra L.	1,726.0	930.6	34,403.6	16,677.6
Stratum total	6,604.0		131,750.4	
Sapling				
Abies balsamea (L.) Mill.	410.9	265.3	3,436.3	2,218.7
Acer rubrum L.	373.4	29.0	6,715.1	524.9
Betula papyrifera Marsh.	32.4	18.7	1,142.9	639.5
Picea glauca (Moench) Voss.	33.1	-	226.8	-
Pinus banksiana Lamb.	14.8	-	162.9	-
Pinus resinosa Ait.	1.7	1.5	15.9	14.3
Pinus strobus L.	14.6	-	260.4	-
Populus grandidentata Michx.	2.7	-	125.2	-
Populus tremuloides Michx.	5.2	3.2	147.5	90.3
Quercus rubra L.	543.2	262.2	5,732.0	2,833.1
Sorbus americana Marsh.	2.9	-	27.5	-
Stratum total	1,434.9		17,992.5	
Tree seedling				
Abies balsamea (L.) Mill.	28.4	11.9	61.5	27.3
Acer rubrum L.	12.3	3.0	124.8	37.3
Betula papyrifera Marsh.	.6	.4	15.5	11.3
Picea glauca (Moench) Voss.	.8	.8	1.2	-
Pinus banksiana Lamb.	.1	-	.9	-
Pinus resinosa Ait.	2.5	-	21.2	-
Pinus strobus L.	4.2	2.9	30.4	22.5
Populus grandidentata Michx.	.5	-	2.8	-
Populus tremuloides Michx.	5.1	3.2	26.7	16.3
Quercus rubra L.	49.6	19.4	280.8	124.9
Sorbus americana Marsh.	3.5	2.1	21.6	13.8
Tilia americana L.	.2	-	.9	-
Stratum total	107.8		588.3	
Tall shrub				
Acer spicatum Lam.	.8	-	1.4	-
Amelanchier Medic. spp.	113.1	109.7	824.2	773.3
Cornus rugosa Lam.	.9	-	1.7	-
Corylus cornuta Marsh.	82.5	23.1	406.0	117.6
Lonicera canadensis Marsh.	1.3	-	4.2	-
Prunus pensylvanica L.F.	<.1	-	.1	-
Salix bebbiana Sarg.	.2	.2	.3	.3
Viburnum rafinesquianum Schult.	.3	-	.7	-
Stratum total	199.2		1,238.6	

(Table 12 continued on next page)

(Table 12.--continued)

Stratum/species	BWC AW MATURE (n = 4)		Total	
	Foliage \bar{X}	SE	Foliage \bar{X}	SE
Low shrub				
<i>Comptonia peregrina</i> B.&B. Small. Rydb.			4.0	2.9
<i>Diervilla Lonicera</i> Mill.			4.2	-
<i>Gaultheria procumbens</i> L.			24.3	9.7
<i>Ribes triste</i> Pall.			<.1	-
<i>Rosa acicularis</i> Lindl.			1.1	.6
<i>Rubus pubescens</i> Raf.			.4	-
<i>Vaccinium angustifolium</i> Ait.			66.1	42.7
<i>Vaccinium myrtilloides</i> Michx.			16.4	6.6
Stratum total			116.6	
Herb				
<i>Anemone quinquefolia</i> L.	<0.1	-		
<i>Aralia nudicaulis</i> L.	15.5	6.6		
<i>Aster ciliolatus</i> Lindl.	.3	-		
<i>Aster macrophyllus</i> L.	74.9	21.6		
<i>Carex</i> L. spp.	7.3	2.1		
<i>Clintonia borealis</i> (Ait.) Raf	5.4	1.3		
<i>Coptis groenlandica</i> (Oeder) Fern	.4	-		
<i>Cornus canadensis</i> L.	5.7	1.6		
<i>Epigaea repens</i> L.	4.8	-		
<i>Fragaria vesca</i> L.	.4	-		
<i>Galium triflorum</i> Michx.	<.1	-		
<i>Goodyera repens</i> (L.) R. Br.	.2	-		
Grasses	30.6	5.4		
<i>Lathyrus ochroleucus</i> Hook.	.4	-		
<i>Maianthemum canadense</i> Desf.	12.0	2.4		
<i>Melampyrum lineare</i> Desr.	5.5	2.6		
<i>Mitella nuda</i> L.	.2	-		
<i>Monotropa uniflora</i> L.	.1	.1		
<i>Pyrola secunda</i> L.	.1	.1		
<i>Streptopus roseus</i> Michx.	1.1	.6		
<i>Trientalis borealis</i> Raf.	.6	.3		
<i>Uvularia sessilifolia</i> L.	.1	.1		
<i>Viola</i> L. spp.	.2	-		
Stratum total	165.9			
Fern and fern allies				
<i>Lycopodium annotinum</i> L.	1.6	1.1		
<i>Lycopodium clavatum</i> L.	2.0	1.8		
<i>Lycopodium obscurum</i> L.	1.7	1.2		
<i>Pteridium aquilinum</i> (L.) Kuhn.	37.7	9.2		
Stratum total	43.0			
Moss				
<i>Calliergonella schreberi</i> (Willd., BR. & Sch.) Grout	31.8	23.4		
<i>Dicranum</i> Hedw. spp.	43.3	18.3		
<i>Polytrichum</i> Hedw. spp.	18.2	10.3		
All other mosses	194.6	50.5		
Stratum total	287.9			
Lichen				
<i>Cladonia rangiferina</i> (L.) Web.	5.3	3.7		
All other lichens	17.2	3.9		
Stratum total	22.5			
Community total			152,205.7	

Table 13.--Aboveground ovendry biomass by species and stratum for the red pine community type of northeastern Minnesota

Stratum/species	(In Kg/ha)			
	BWC AW		MATURE (n = 6)	
	Foliage	Total	X	SE
Tree				
<i>Abies balsamea</i> (L.) Mill.	31.9	17.4	257.6	141.6
<i>Acer rubrum</i> L.	90.9	39.0	3,741.3	1,560.0
<i>Betula papyrifera</i> Marsh	109.3	51.4	3,142.8	1,467.3
<i>Picea glauca</i> (Moench) Voss.	17.4	-	149.4	-
<i>Picea mariana</i> (Mill.) B.S.P.	177.6	91.5	2,074.6	1,317.3
<i>Pinus banksiana</i> Lamb.	1,114.7	814.0	21,922.7	14,230.8
<i>Pinus resinosa</i> Ait.	7,219.0	4,657.2	120,105.5	82,315.4
<i>Pinus strobus</i> L.	3,763.2	1,307.4	84,058.1	29,273.9
<i>Populus grandidentata</i> Michx.	10.1	-	531.5	-
<i>Populus tremuloides</i> Michx.	.8	-	34.2	-
<i>Quercus rubra</i> L.	47.2	27.4	760.6	462.3
<i>Thuja occidentalis</i> L.	34.8	-	294.9	-
Stratum total	12,616.9		237,073.2	
Sapling				
<i>Abies balsamea</i> (L.) Mill.	12.9	8.4	109.2	70.9
<i>Acer rubrum</i> L.	82.0	51.9	1,266.1	844.7
<i>Betula papyrifera</i> Marsh.	15.7	12.3	631.5	495.4
<i>Picea glauca</i> (Moench) Voss.	2.7	1.8	17.2	10.9
<i>Picea mariana</i> (Mill.) B.S.P.	52.5	30.7	291.1	177.1
<i>Pinus banksiana</i> Lamb.	8.8	6.5	85.6	62.3
<i>Pinus resinosa</i> Ait.	22.2	14.1	219.2	139.9
<i>Pinus strobus</i> L.	36.6	16.4	651.0	293.5
<i>Quercus rubra</i> L.	34.1	18.4	325.2	177.4
<i>Thuja occidentalis</i> L.	12.2	-	60.7	-
Stratum total	279.7		3,656.8	
Tree seedling				
<i>Abies balsamea</i> (L.) Mill.	2.1	1.7	4.5	3.6
<i>Acer rubrum</i> L.	7.2	3.9	96.9	47.8
<i>Betula papyrifera</i> Marsh.	.6	.4	11.9	9.0
<i>Picea glauca</i> (Moench) Voss.	6.1	4.4	12.0	9.2
<i>Picea mariana</i> (Mill.) B.S.P.	34.0	18.1	63.0	32.1
<i>Pinus resinosa</i> Ait.	1.8	1.1	13.3	8.4
<i>Pinus strobus</i> L.	21.6	9.4	158.0	69.2
<i>Populus tremuloides</i> Michx.	.8	.3	4.4	1.7
<i>Quercus rubra</i> L.	13.2	5.9	66.4	31.3
<i>Sorbus americana</i> Marsh.	.4	.3	2.2	1.5
Stratum total	87.8		432.6	
Tall shrub				
<i>Alnus crispa</i> (Ait.) Pursh.	10.7	5.0	77.5	36.0
<i>Amelanchier</i> Medic. spp.	5.1	3.8	56.0	38.8
<i>Corylus cornuta</i> Marsh.	29.8	18.0	156.8	95.6
<i>Lonicera canadensis</i> Marsh.	.1	-	.2	-
<i>Salix bebbiana</i> Sarg.	.7	.5	2.5	2.4
Stratum total	46.4		293.0	
Low shrub				
<i>Arctostaphylos Uva-ursi</i> (L.) Spreng.			5.8	-
<i>Chimaphila umbellata</i> (L.) Bart.			1.0	.7
<i>Comptonia peregrina</i> B.&B. Small. Rydb.			.2	.2
<i>Diervilla</i> Lonicera Mill.			2.0	1.0
<i>Gaultheria procumbens</i> L.			33.2	13.1
<i>Juniperus communis</i> L.			89.0	59.7
<i>Vaccinium angustifolium</i> Ait.			150.6	53.9
<i>Vaccinium myrtilloides</i> Michx.			53.7	25.3
Stratum total			335.5	

(Table 13 continued on next page)

(Table 13.--continued)

Stratum/species	BWC AW MATURE (n = 6)		Total	
	Foliage \bar{X}	SE	\bar{X}	SE
Anemone <u>quinquefolia</u> L.	<0.1	-		
Aralia <u>nudicaulis</u> L.	6.0	3.9		
Aster <u>macrophyllus</u> L.	38.9	22.7		
Carex L. spp.	4.5	3.2		
Clintonia <u>borealis</u> (Ait.) Raf.	1.7	1.4		
Cornus <u>canadensis</u> L.	7.1	3.3		
Goodyera <u>repens</u> (L.) R. Br.	.1	.1		
Goodyera <u>tesselata</u> Lodd.	.2	-		
Grasses	32.8	21.2		
Linnaea <u>borealis</u> L.	.5	-		
Maianthemum <u>canadense</u> Desf.	6.8	1.0		
Melampyrum <u>lineare</u> Desr.	6.0	2.2		
Streptopus <u>roseus</u> Michx.	.1	-		
Trientalis <u>borealis</u> Raf.	.2	.1		
Stratum total	105.0			
Fern and fern allies				
Lycopodium <u>annotinum</u> L.	.6	-		
Lycopodium <u>clavatum</u> L.	.9	.6		
Lycopodium <u>obscurum</u> L.	1.0	-		
Polypodium <u>virginianum</u> L.	.1	-		
Pteridium <u>aquilinum</u> (L.) Kuhn.	10.5	6.8		
Stratum total	13.1			
Moss				
Calliergonella <u>schreberi</u> (Willd., BR. & Sch.) Grout	788.5	207.5		
Dicranum Hedw. spp.	246.6	62.8		
Hypnum <u>crista-castrensis</u> Hedw.	.7	-		
Polytrichum Hedw. spp.	36.8	12.0		
Sphagnum L. spp.	16.1	-		
All other mosses	160.2	39.7		
Stratum total	1,248.9			
Lichen				
Cladonia <u>mitis</u> Sandst.	122.6	83.9		
Cladonia <u>rangiferina</u> (L.) Web	364.2	205.6		
Cladonia <u>sylvatica</u> (L.) Hoffm.	66.1	-		
All other lichens	62.2	36.8		
Stratum total	615.1			
Community total			243,773.2	

Table 14.--Aboveground ovendry biomass by species and stratum for the jack pine-fir community type of northeastern Minnesota.

(Table 14 continued on next page)

Table 14.--continued)

	Prayer Lake Fire						Little Sioux Fire						1-5 n = 2					
	1 n = 2			1 n = 2			3 n = 2			5 n = 2			1-5 n = 2					
	Foliage	\bar{X}	SE	Foliage	\bar{X}	SE	Foliage	\bar{X}	SE	Foliage	\bar{X}	SE	Foliage	\bar{X}	SE			
Herbs																		
Actaea rubra (Ait.) Willd.																		
Anemone quinquefolia L.				<0.1	-		0.2	-					<0.1	-				
Antennaria spp. Gaertn.	0.1	-		.5	-								5.0	-				
Apocynum androsaemifolium L.				.2	.1		26.2	-										
Aralia hispida Vent.	.1	-		6.4	4.2		39.3	6.7										
Aralia nudicaulis L.																		
Aster ciliolatus Lindl.																		
Aster macrophyllus L.	75.6	-		289.4	3.1		284.6	17.8										
Zarex L. spp.	14.2	7.2		74.2	28.4		183.0	122.6										
Cytisus alpinus L.																		
Spiraea betulifolia (Ait.) Raf.																		
Spiraea prunifolia (Oeder) Fern.																		
Zonalornis maculata Raf.																		
Cornus canadensis L.																		
Cardaliss semperfervens (L.) Pers.	18.2	17.2		17.6	13.4		71.1	35.9										
Epilobium angustifolium L.	8.6	1.2					4.5	-										
Fragaria vesca L.																		
Fragaria Virginiana Duchesne.																		
Gaultheria triflorum Michx.																		
Geranium maculatum L.																		
Gosoyer repens (L.) R. Br.	197.9	80.6	197.9	80.6	13.6	-	<.1	-										
Gosoyer tessellata Lodd.																		
Grasses																		
Zizaniopsis miliacea L.																		
Zizaniopsis miliacea Hook.																		
Zizaniopsis miliacea L.																		
Zizaniopsis venosa Muhl.																		
Zizaniopsis ... spp.																		
Linnæa borealis L.	<.1	-		1.2	.5		40.4	34.6										
Lycopodium uniflorum																		
Melanthium canadense Desf.	.1	0		3.0	2.2		32.6	10.4										

	Prayer Lake Fire						Little Sioux Fire						1-5 n = 2					
	1 n = 2			1 n = 2			3 n = 2			5 n = 2			1-5 n = 2					
	Foliage	\bar{X}	SE	Foliage	\bar{X}	SE	Foliage	\bar{X}	SE	Foliage	\bar{X}	SE	Foliage	\bar{X}	SE			
Herbs (cont.)																		
Monotropa uniflora Desr.							0.2	-										
Monotropa uniflora L.		3.0	-															
Moneses uniflora (L.) Gray																		
Petasites palmatus (Ait.) Gray																		
Polygonia caerulea Willd.																		
Polygonum citrinum Michx.	7.8	7.1		2.2	-		15.8	13.2										
Pyrola asarifolia Michx.																		
Pyrola elliptica Nutt.																		
Pyrola secunda L.																		
Pyrola viridis Schleid.																		
Pyrola L. spp.																		
Solidago uliginosa																		
Streptopus roseus Michx.																		
Tephroseris integrifolia Spreng.																		
Trifolium bifoliatum Raf.				.5	-													
Trifolium repens																		
Vicia americana Muhl.																		
Viola L. spp.																		
Waldsteinia fragarioides (Michx.) Tratt.																		
Fern and fern allies																		
Athyrium filix-femina (L.) Roth.																		
Dryopteris Phegopteris L.																		
Dryopteris spinulosa (O.F. Muell.) Watt																		
Equisetum L. spp.																		
Gymnocarpium Dryopteris (L.) Newm.																		
Lycopodium acutulum L.																		
Lycopodium clavatum L.																		
Lycopodium complanatum L.				.1	-		23.8	-										
Stratum total																		

	Prayer Lake Fire						Little Sioux Fire						1-5 n = 2					
	1 n = 2			1 n = 2			3 n = 2			5 n = 2			1-5 n = 2					
	Foliage	\bar{X}	SE	Foliage	\bar{X}	SE	Foliage	\bar{X}	SE	Foliage	\bar{X}	SE	Foliage	\bar{X}	SE			
Fern and fern allies (cont.)																		
Lycopodium obscurum L.																		
Osmunda claytoniana L.																		
Polyodium virginianum L.																		
Pteridium aquilinum (L.) Kuhn.																		
Woodia ilvensis (L.) R. Br.																		
Stratum total				203.1	136.1		261.1	56.4										
Calliergonella schreberi (Willd., Br. & Son.) Grout																		
Hypoleium splendens (Hedw.) B.S.G.																		
Hypnum crista-caeruleum Hedw.																		
Polytrichum Hedw. spp.	2.3	1.9																
Spiranthes L. spp.																		
All other mosses																		
Stratum total																		
Lichen																		
Cladonia mitis Sandst.																		
Cladonia rangiferina (L.) Web.																		
Cladonia sylvatica (L.) Hoffm.																		
All other lichens																		
Stratum total																		
Unidentified plants	.1	0		<.1	-													
Community total				354.4			1,202.2											

Isabella Plantations										Cherokee-Frost Lake										BWCAN MATURE n = 10			
6-10 n = 2		16-20 n = 2		21-25 n = 2		26-30 n = 2																	
Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total	
\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE		
<0.1		<0.1		<0.1		<0.1		0.1		0.1		0.5		0.4		1.7		1.6					
<.1		<.1		<.1		<.1		.1		.1		.5		-		.1		-					
<0.1		<.1		<.1		<.1		.1		.1		.1		.1		.1		.1		.1			
<.1		<.1		<.1		<.1		.2		.1		.1		.1		.1		.1		.1			
<.1		<.1		<.1		<.1		.4		.3		.3		.2		.2		.2		.2			
<.1		<.1		<.1		<.1		.1		.1		.1		.1		.1		.1		.1			
<.1		<.1		<.1		<.1		.6		.2		.4		.2		.4		.2		.2			
<.1		<.1		<.1		<.1		.1		.1		.1		.1		.1		.1		.1			
1.3		1.3		1.3		1.3		.8		.4		.7		.5		.7		.3		.3			
312.1		369.7		436.6		197.8		198.5		198.5		237.5		237.5		.8		-					
<.1		<.1		<.1		<.1		.5		.1		1.6		1.0		.1		-					
<.1		<.1		<.1		<.1		.3		.2		.2		.2		2.1		1.9					
<.1		<.1		<.1		<.1		1.1		.1		3.9		2.1		4.3		2.2					
<.1		<.1		<.1		<.1		0		-		.3		-		-							

Table 15.--Aboveground ovendry biomass by species and stratum for the jack pine-black spruce community type of northeastern Minnesota

Tree	(In Kg/ha)																									
	Little Sioux Fire						6-10						11-15													
	Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total			
	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE		
Abies balsamea (L.) Mill.																										
Acer rubrum L.																										
Betula papyrifera Marsh.																										
Picea glauca (Moench) Voss.																										
Picea mariana (Mill.) B.S.P.																										
Pinus banksiana Lamb.																										
Pinus resinosa Ait.																										
Pinus strobus L.																										
Populus grandidentata Michx.																										
Populus tremuloides Michx.																										
Quercus rubra L.																										
Thuja occidentalis L.																										
Stratum total																										
Sapling																										
Abies balsamea (L.) Mill.																										
Acer rubrum L.																										
Betula papyrifera Marsh.																										
Picea glauca (Moench) Voss.																										
Picea mariana (Mill.) B.S.P.																										
Pinus banksiana Lamb.																										
Pinus resinosa Ait.																										
Pinus strobus L.																										
Populus grandidentata Michx.																										
Populus tremuloides Michx.																										
Quercus rubra L.																										
Sorbus americana Marsh.																										
Thuja occidentalis L.																										
Stratum total																										

Tree seedling*	(In Kg/ha)																										
	Little Sioux Fire						6-10						11-15														
	Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total				
	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE			
Abies balsamea (L.) Mill.																											
Acer rubrum L.																											
Betula papyrifera Marsh.																											
Picea glauca (Moench) Voss.																											
Picea mariana (Mill.) B.S.P.	1.4		60.7		<0.1		95.8		658.4		1,387.8																
Pinus banksiana Lamb.																											
Pinus resinosa Ait.																											
Pinus strobus L.																											
Populus grandidentata Michx.																											
Populus tremuloides Michx.	34.9		13.9		1.4		34.2		53.0		235.8		31.5		15.4		227.0		108.4		24.1		18.4		183.8		134.5
Quercus rubra L.																											
Sorbus americana Marsh.																											
Thuja occidentalis L.																											
Stratum total	36.3		75.5		131.5		1,309.1		4,437.5		35.5		289.0		24.6		-										
Tall shrub																											
Acer spicatum Lam.																											
Ailanthus altissima (L.) Pursh.																											
Ailanthus rugosissima (DuRoi) Spreng.																											
Amelanchier spp.	14.5																										
Cornus rugosa L.																											
Cornus stolonifera Michx.																											
Dryopteris marginalis Marsh.																											
Lonicera canadensis Marsh.																											
Lonicera dioica L.																											
Lonicera hirsutissima Nutt.																											
Gaultheria procumbens L.																											
Juniperus communis L.																											
Ledum groenlandicum Oeder.																											
Ribes spp.																											
Rosa blanda L.																											
Rosa acicularis Lindl.																											
Rubus pensylvanicus Nutt.																											
Rubus pubescens Raf.																											
Rubus strigosus Michx.																											
Vaccinium angustifolium Ait.	12.0																										
Vaccinium myrtillifolium Michx.	9.6																										
Stratum total	71.4																										

Isabella Plantations										Cherokee-Frost Lake										BWCW MATURE n = 17						
16-20 n = 4				21-25 n = 3				26-30 n = 6				33 n = 3														
Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total				
\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE			
136.5	94.4	988.4	681.8	1,008.7	884.3	21,386.5	17,343.0	204.6	97.1	5,396.9	2,235.2	307.0	293.6	12,681.3	11,989.5	52.2	32.4	416.6	259.7	28.9	13.3	1,021.3	456.0			
339.5	-	2,816.6	-	262.8	-	2,114.3	-	34.8	-	288.0	-	72.7	33.9	1,993.6	976.2	70.1	36.9	545.4	272.4	416.2	266.5	12,463.3	7,813.0			
5,502.4	779.1	57,614.4	8,610.8	416.0	356.6	2,962.8	2,665.1	7.5	-	41.5	-	92.9	-	466.2	-	3,891.9	664.9	30,865.7	5,364.6	496.9	290.4	6,726.8	3,956.4			
285.1	235.4	7,361.0	6,232.5	294.1	10,595.0	5,549.3	3,598.0	1,595.8	58,030.5	26,973.2	8,631.0	4,949.8	101,547.7	50,134.3	5,401.6	701.2	101,136.5	12,660.6	147.2	-	2,616.0	-	10.1	7.3	773.5	565.4
6,904.7	81,243.7	6,154.2	-	1,178.1	-	1,008.7	-	235.5	218.1	8,322.1	7,780.9	102.9	-	4,004.2	-	127.2	29.8	6,069.6	1,458.8	8.4	-	156.5	-	10,339.6	153,844.2	
6,904.7	81,243.7	6,154.2	-	1,178.1	-	1,008.7	-	8,705.9	126,939.7	9,444.7	-	122,717.8	-	10,339.6	-	36.1	15.6	307.6	133.0	10.0	4.9	177.4	95.2			
133.9	-	-	-	10.3	-	-	-	133.9	-	5,115.1	-	29.7	9.0	1,109.9	338.7	10.3	-	60.1	-	15.7	10.2	100.5	65.1			
234.6	-	-	-	854.0	-	-	-	234.6	-	1,087.0	-	1,751.9	339.9	10,259.3	1,974.8	854.0	-	8,143.9	-	70.5	34.2	750.4	371.7			
.2	.2	.2	.2	.2	.2	.2	.2	.2	.2	.2	.2	.2	.2	.2	.2	.2	.2	.2	.2	.2	.2	.2	.2			
1,233.0	14,406.7	1,937.6	-	1,233.0	14,406.7	1,937.6	-	1,233.0	14,406.7	1,937.6	-	1,233.0	14,406.7	1,937.6	-	1,233.0	14,406.7	1,937.6	-	1,233.0	14,406.7	1,937.6	-			

Isabella Plantations										Cherokee-Frost Lake										BWCW MATURE n = 17					
16-20 n = 4				21-25 n = 3				26-30 n = 6				33 n = 3													
Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total			
\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE		
0.2	0.1	1.1	0.8	16.2	-	0.1	0.1	0.2	0.2	9.3	1.8	246.7	44.4	5.6	2.2	98.7	40.9	3.0	-	5.1	-	1.2	-	13.1	5.9
11.2	8.4	65.6	43.4	1.1	1.0	11.5	9.7	.1	-	1.3	-	.7	-	4.3	-	2.7	1.1	20.7	9.0	1.1	-	.1	-	4.1	2.0
11.4	-	66.7	7.3	63.1	-	7.2	-	1.5	-	125.0	-	502.9	-	51.1	-	2.2	1.1	20.7	9.0	1.1	-	.2	-	4.1	2.0
98.5	69.4	965.5	684.9	180.9	111.8	117.2	63.5	10.3	-	95.6	-	2.2	2.1	9.0	8.7	1.0	.8	8.2	7.9	6.1	4.1	48.2	32.3		
38.0	13.6	320.1	132.1	25.1	17.2	128.2	-	29.1	18.2	279.2	196.2	<.1	<.1	.7	.1	.4	.2	6.3	3.1	1.3	.7	8.0	4.8		
97.4	20.3	889.2	178.4	276.0	183.5	2,304.2	1,515.0	246.8	42.6	2,092.5	351.8	33.2	32.8	200.0	199.6	15.3	10.7	90.8	65.9	1.4	1.4	14.4	8.8		
2.5	-	.8	-	<.1	-	.1	-	.5	-	4.8	-	.6	.4	2.3	1.5	.5	-	.1	.1	.1	.1	.1	.1		
9.9	-	25.8	-	2.6	-	26.1	-	1.1	.5	11.9	5.2	4.8	4.8	36.3	36.0	<.1	<.1	<.1	<.1	2.6	-	3.1	1.1		
89.1	-	-	-	3.1	2.6	17.8	15.9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		

Isabella Plantations										Cherokee-Frost Lake										BWCW MATURE n = 17					
16-20 n = 4				21-25 n = 3				26-30 n = 6				33 n = 3													
Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total			
\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE		
248.6	B,304.9	518.0	4,763.6	5.2	3.1	36.1	15.2	1.3	-	8.0	-	0.1	0.1	0.2	0.1	2.3	1.9	28.6	26.8	.6	-	2.1	1.5	.3	.1
74.4	33.4	68.0	31.7	10.2	5.0	9.7	5.1	14.6	5.0	8.1	5.2	10.2	6	10.2	4.7	10.2	6	4.7	4	8.1	5.2	10.2	4.7	10.5	3.4
5.7	-	8.0	5.4	19.3	11.2	3.6	1.3	4.9	-	.8	-	.8	.8	.8	-	36.2	27.3	1.2	-	.8	-	25.4	9.8	4.0	1.9
.2	.7	22.3	10.6	5.6	3.0	8.0	2.2	2.2	-	2.2	-	1.2	-	1.2	-	1.2	.9	1.2	.9	1.2	.9	1.8	1.6	.1	.1
1.2	.8	25.2	9.0	10.2	5.0	9.7	5.1	.4	.2	.3	.3	.3	.3	.3	.3	25.2	27.3	.3	.3	.3	.3	25.4	9.8	4.0	1.9
20.8	11.4	19.3	11.2	3.6	1.3	3.6	1.3	.8	-	.8	-	.8	.8	.8	-	32.6	37.3	11.2	11.2	11.2	11.2	32.6	37.3	11.2	11.2
32.6	12.5	120.9	9.5	8.3	-	42.4	-	23.0	-	23.0	-	23.0	-	23.0	-	32.6	37.3	9.5	9.5	9.5	9.5	32.6	37.3	9.5	9.5
189.4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

(Table 15 continued on next page)

(Table 15.--continued)

	Little Sioux Fire										11-15									
	1 n = 1		3 n = 1		5 n = 1		6-10 n = 6		11-15 n = 4											
	Foliage	Total	Foliage	Total	Foliage	Total	Foliage	Total	Foliage	Total										
	X	SE	X	SE	X	SE	X	SE	X	SE										
Herb																				
<i>Actaea rubra</i> (Ait.) Willd.							<0.1	-												
<i>Anaphalis margaritacea</i> (L.) C.B. Clarke							<.1	-												
<i>Anemone quinquefolia</i> L.					0.2		<.1	<0.1												
<i>Antennaria neodioica</i> Greene																				
<i>Antennaria</i> spp. Gaertn.							<.1	<.1												
<i>Apocynum androsaemifolium</i> L.							1.4	.8												
<i>Aralia hispida</i> Vent.	.7																			
<i>Aralia nudicaulis</i> L.	1.4		383.6		55.1		3.8	1.4	11.0	6.5										
<i>Aster ciliolatus</i> Lindl.			21.9		26.6		.7	-												
<i>Aster macrophyllus</i> L.	88.3				184.1		45.7	30.1	87.6	49.3										
<i>Carex</i> spp.			278.6		124.1		<.1	<.1	11.4	5.8										
<i>Clintonia borealis</i> (Ait.) Raf.	8.8		12.7		28.9		8.1	5.4	<.1	<.1										
<i>Coptis groenlandica</i> (Oeder) Fern.																				
<i>Cornus canadensis</i> L.																				
<i>Epilobium angustifolium</i> L.	41.4		227.7		255.9		9.5	5.2	12.9	2.8										
<i>Fragaria vesca</i> L.							2.9	1.3	.4	.2										
<i>Fragaria virginiana</i> Duchesne.							8.1	3.1	3.2	.7										
<i>Gallium triflorum</i> Michx.							<.1	<.1												
<i>Geranium bicknellii</i> Britt.	<.1		6.6																	
<i>Goodyera repens</i> (L.) R. Br.																				
<i>Goodyera tesselata</i> Lodder.																				
Grasses	2.4		21.4		10.4		158.0	61.5	115.2	29.7										
<i>Habenaria orbiculata</i> (Pursh) Torr.																				
<i>Hepatica americana</i> (D.C.) Ker.							<.1	-												
<i>Lactuca</i> L. spp.							<.1	-												
<i>Lathyrus ochroleucus</i> Hook.							4.1	2.0	.7	-										
<i>Lathyrus palustris</i> L.							.2	-												
<i>Lathyrus venosus</i> Muhl.							.2	-												
<i>Linnea borealis</i> L.							1.9	1.0	17.0	8.0										
<i>Maianthemum canadense</i> Desf.	5.4		34.1		.9		2.4	1.0	5.8	1.9										
Grasses					11.3		2.4		<.1	-										

	Little Sioux Fire										11-15									
	1 n = 1		3 n = 1		5 n = 1		6-10 n = 6		11-15 n = 4											
	Foliage	Total	Foliage	Total	Foliage	Total	Foliage	Total	Foliage	Total										
	X	SE	X	SE	X	SE	X	SE	X	SE										
Herb (cont.)																				
<i>Monotropa uniflora</i> L.							<0.1	-												
<i>Petasites palmatus</i> (Ait.) Gray							<.1	<0.1												
<i>Petasites vitifolius</i> Rydb.									<.1	<.1										
<i>Polygonia paucifolia</i> Willd.																				
<i>Polygonum cilinode</i> Michx.																				
<i>Potentilla tridentata</i> Soland.	0.9		25.8		2.0															
<i>Pyrola elliptica</i> Nutt.																				
<i>Pyrola rotundifolia</i> L.																				
<i>Pyrola secunda</i> L.																				
<i>Pyrola virens</i> Schweigg.																				
<i>Pyrola</i> L. spp.																				
<i>Sedum</i> L. spp.																				
<i>Solidago</i> L. spp.																				
<i>Streptopus roseus</i> Michx.																				
<i>Taraxacum officinale</i> Web.																				
<i>Trientalis borealis</i> Raf.	.2																			
<i>Vicia americana</i> Muhl.																				
<i>Viola adunca</i> J. E. Smith																				
<i>Viola</i> L. spp.																				
<i>Walsteinia fragarioides</i> (Michx.) Tratt.																				
Stratum total	149.6		1,136.7		580.5		249.7		268.4											
Fern and fern allies																				
<i>Dryopteris spinulosa</i> (O.F. Muell.) Watt																				
<i>Lycopodium annotinum</i> L.																				
<i>Lycopodium clavatum</i> L.																				
<i>Lycopodium complanatum</i> L.																				
<i>Lycopodium lucidulum</i> Michx.																				
<i>Lycopodium obscurum</i> L.																				
<i>Polyodium virginianum</i> L.																				
<i>Pteridium aquilinum</i> (L.) Kuhn.	273.7		403.0		599.8		11.1	8.2	11.6	6.8										
Stratum total	273.7		437.1		599.8		11.5		11.8											

	Little Sioux Fire										11-15									
	1 n = 1		3 n = 1		5 n = 1		6-10 n = 6		11-15 n = 4											
	Foliage	Total	Foliage	Total	Foliage	Total	Foliage	Total	Foliage	Total										
	X	SE	X	SE	X	SE	X	SE	X	SE										
Moss																				
<i>Calliergonella schreber</i>																				
(WILLD., BR. & SCH.) Grout																				
<i>Dicranum</i> Hedw. spp.																				
<i>Hyalacomium splendens</i> (Hedw.) B.S.G.																				
<i>Hypnum cristatum</i> (Hedw.) Castrovilli																				
<i>Polytrichum</i> Hedw. spp.																				
All other mosses																				
Stratum total			12.3				10.2													
Lichens																				
<i>Cladonia alpestris</i> (L.) Rabenh.																				
<i>Cladonia mitis</i> Sandst.																				
<i>Cladonia rangiferina</i> (L.) Web.																				
<i>Cladonia sylvatica</i> (L.) Hoffm.																				
<i>Peltigera malacea</i> (L.) Willd.																				
All other lichens																				
Stratum total			12.3				10.2													
Liverworts																				
Unidentified																				
Community total			.1				<.1													
			546.1		2,563.5		7,441.2		74,349.9											
										45,501.6										

Isabella Plantations										Cherokee-Frost Lake										BWC AW MATURE n = 17	
16-20 n = 4				21-25 n = 3				26-30 n = 6				33 n = 3									
Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total			
\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE
<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	0.2	-	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
.1	-	.1	-	.1	-	.1	-	2.8	-	.2	.2	.2	.2	.2	.2	.2	.2	.2	.2	.2	.2
.6	-	.6	-	.6	-	.6	-	.8	-	.8	-	.8	-	.8	-	.8	-	.8	-	.8	-
7.2	5.5	13.7	7.7	11.5	6.9	15.9	8.8	4.5	1.6	4.5	1.6	4.5	1.6	4.5	1.6	4.5	1.6	4.5	1.6	4.5	1.6
52.5	14.0	99.8	38.1	40.8	15.4	78.0	59.2	35.5	7.3	35.5	7.3	35.5	7.3	35.5	7.3	35.5	7.3	35.5	7.3	35.5	7.3
<.1	-	<.1	-	<.1	-	.9	-	2.1	1.3	2.1	1.3	2.1	1.3	2.1	1.3	2.1	1.3	2.1	1.3	2.1	1.3
7.0	.2	22.8	11.1	4.4	1.5	3.2	1.4	3.1	1.3	3.1	1.3	3.1	1.3	3.1	1.3	3.1	1.3	3.1	1.3	3.1	1.3
.1	-	<.1	-	<.1	-	.2	-	.2	-	.2	-	.2	-	.2	-	.2	-	.2	-	.2	-
5.8	3.8	16.6	6.9	9.6	3.6	41.2	28.7	25.4	8.0	25.4	8.0	25.4	8.0	25.4	8.0	25.4	8.0	25.4	8.0	25.4	8.0
1.8	-	4.1	-	4.1	3.4	.6	-	.6	-	.6	-	.6	-	.6	-	.6	-	.6	-	.6	-
2.8	1.4	2.2	-	.8	.5	1.5	1.0	.1	-	.1	-	.1	-	.1	-	.1	-	.1	-	.1	-
165.0	93.7	134.1	115.7	47.9	8.0	9.0	5.3	8.6	2.4	8.6	2.4	8.6	2.4	8.6	2.4	8.6	2.4	8.6	2.4	8.6	2.4
.9	-	.7	-	.7	-	.7	-	.3	-	.3	-	.3	-	.3	-	.3	-	.3	-	.3	-
5.8	4.3	5.8	4.7	10.9	4.9	29.5	25.1	12.7	4.4	12.7	4.4	12.7	4.4	12.7	4.4	12.7	4.4	12.7	4.4	12.7	4.4
2.9	1.5	4.1	1.8	5.3	1.4	9.5	6.8	9.9	1.7	9.9	1.7	9.9	1.7	9.9	1.7	9.9	1.7	9.9	1.7	9.9	1.7
<.1	-	-	-	-	-	-	-	.4	-	.4	-	.4	-	.4	-	.4	-	.4	-	.4	-

Isabella Plantations										Cherokee-Frost Lake										BWC AW MATURE n = 17	
16-20 n = 4				21-25 n = 3				26-30 n = 6				33 n = 3									
Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total			
\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE
<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1
2.7	.4	2.6	1.5	0.6	-	0.6	-	0.6	-	0.6	-	0.6	-	0.6	-	0.6	-	0.6	-	0.6	-
.7	-	.8	-	.8	-	.8	-	.8	-	.8	-	.8	-	.8	-	.8	-	.8	-	.8	-
.1	-	.1	-	.1	-	.1	-	.1	-	.1	-	.1	-	.1	-	.1	-	.1	-	.1	-
.3	-	.4	-	.2	-	.2	-	.2	-	.2	-	.2	-	.2	-	.2	-	.2	-	.2	-
255.3	304.7	139.2	198.7	105.3	1.0	14.5	12.8	3.7	1.8	3.7	1.8	3.7	1.8	3.7	1.8	3.7	1.8	3.7	1.8	3.7	1.8
<.1	-	<.1	0	<.1	-	<.1	-	<.1	-	<.1	-	<.1	-	<.1	-	<.1	-	<.1	-	<.1	-
<.1	-	<.1	<.1	2.2	-	1.1	-	1.1	-	1.1	-	1.1	-	1.1	-	1.1	-	1.1	-	1.1	-
49.1	28.4	77.4	-	28.1	22.8	15.6	-	15.6	-	15.6	-	15.6	-	15.6	-	15.6	-	15.6	-	15.6	-
49.2	77.5	77.5	-	30.5	-	22.8	-	22.8	-	22.8	-	22.8	-	22.8	-	22.8	-	22.8	-	22.8	-

Isabella Plantations										Cherokee-Frost Lake										BWC AW MATURE n = 17		
16-20 n = 4				21-25 n = 3				26-30 n = 6				33 n = 3										
Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total		Foliage		Total				
\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	\bar{X}	SE	
386.5	241.1	1,920.9	239.8	25.1	9.5	212.3	32.2	125.3	77.2	25.8	15.4	125.3	77.2	17.3	6.9	198.4	47.4	673.0	2,395.2	66.9	-	
44.1	17.6	113.3	63.9	.9	-	20.5	12.4	7.1	1.8	2.5	-	7.1	1.8	2.5	-	1.6	-	111.5	34.1	15.1	-	
60.4	26.9	218.3	-	60.4	26.9	111.5	-	34.1	15.1	34.1	15.1	34.1	15.1	34.1	15.1	34.1	15.1	34.1	15.1	34.1	15.1	
84,109.2	105,064.5	130,223.2	135,896.6	170,148.0																		

Table 16.--Aboveground ovendry biomass by species and stratum for the black spruce-feather moss community type of northeastern Minnesota

Tree	(in Kg/ha)												BNGAW			MATURE		
	11-15 n = 3			Isabelia Plantations n = 3			16-20 n = 3			21-25 n = 1			Foliage			n = 19		
	Foliage X	SE	Total X	Foliage X	SE	Total X	Foliage X	SE	Total X	Foliage X	SE	Foliage X	SE	Total X	SE	Total X	SE	
<i>Abies balsamea</i> (L.) Mill.	5.9	-	66.2	-	62.0	56.6	874.7	810.4	1,297.6	31,235.0	-	871.8	298.7	6,946.4	2,382.7	-	-	
<i>Acer rubrum</i> L.	-	-	-	-	-	-	-	-	-	-	-	1.1	-	45.4	-	-	-	
<i>Betula papyrifera</i> Marsh.	1.3	-	7.4	335.0	-	2,332.9	-	-	-	-	-	396.5	158.1	9,494.7	3,540.5	-	-	
<i>Picea glauca</i> (Moench) Voss.	99.2	72.5	864.1	690.9	-	-	-	-	-	-	-	455.3	159.1	3,986.4	1,365.0	-	-	
<i>Picea mariana</i> (Mill.) B.S.P.	94.2	72.1	794.5	634.1	13.9	-	-	-	-	-	-	5,642.7	1,115.0	46,118.7	9,068.4	-	-	
<i>Pinus banksiana</i> Lamb.	1,648.5	650.3	16,669.2	6,699.7	1,365.2	853.3	14,681.7	9,041.1	1,149.6	6,235.0	12,950.1	1,292.8	489.0	24,828.0	9,159.2	-	-	
<i>Pinus resinosa</i> Ait.	-	-	-	-	-	-	-	-	-	-	-	188.6	139.4	2,451.7	1,804.4	-	-	
<i>Pinus strobus</i> L.	-	-	-	-	-	-	-	-	-	-	-	207.7	98.1	4,528.7	2,121.4	-	-	
<i>Populus grandidentata</i> Michx.	237.2	82.6	4,757.4	1,720.1	102.6	71.8	2,086.5	1,445.0	-	-	-	8.6	7.1	624.2	534.1	-	-	
<i>Populus tremuloides</i> Michx.	-	-	-	-	-	-	-	-	-	-	-	405.9	121.3	19,356.7	5,898.3	-	-	
<i>Sorbus americana</i> Marsh.	-	-	-	-	-	-	-	-	-	-	-	-.8	-.8	8.9	-.9	-	-	
<i>Thuja occidentalis</i> L.	-	-	-	-	-	-	-	-	-	-	-	41.7	76.1	554.8	301.9	-	-	
Stratum total	2,086.3	23,158.8	5,440.5	60,021.9	3,189.4	52,449.7	9,547.9	9,547.9	9,547.9	118,944.6	118,944.6	2,542.8	19,187.2	19,187.2	19,187.2	19,187.2	19,187.2	19,187.2
Sapling	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Abies balsamea</i> (L.) Mill.	-	-	-	-	-	-	-	-	-	-	-	465.1	188.0	3,922.6	1,583.3	-	-	
<i>Acer rubrum</i> L.	-	-	-	-	-	-	-	-	-	-	-	2.0	1.4	33.5	24.5	-	-	
<i>Betula papyrifera</i> Marsh.	-	-	-	-	-	-	-	-	-	-	-	65.2	16.3	2,416.4	605.0	-	-	
<i>Picea glauca</i> (Moench) Voss.	-	-	-	-	-	-	-	-	-	-	-	13.5	7.2	84.8	44.9	-	-	
<i>Picea mariana</i> (Mill.) B.S.P.	-	-	-	-	-	-	-	-	-	-	-	1,894.4	532.1	11,215.9	3,144.5	-	-	
<i>Pinus banksiana</i> Lamb.	-	-	-	-	-	-	-	-	-	-	-	25.9	16.1	265.6	167.5	-	-	
<i>Pinus resinosa</i> Ait.	-	-	-	-	-	-	-	-	-	-	-	1.1	1.7	10.2	7.0	-	-	
<i>Pinus strobus</i> L.	-	-	-	-	-	-	-	-	-	-	-	3.6	2.6	62.3	44.9	-	-	
<i>Populus grandidentata</i> Michx.	-	-	-	-	-	-	-	-	-	-	-	1.4	1.2	57.6	49.7	-	-	
<i>Populus tremuloides</i> Michx.	-	-	-	-	-	-	-	-	-	-	-	36.2	11.7	954.8	311.6	-	-	
<i>Sorbus americana</i> Marsh.	-	-	-	-	-	-	-	-	-	-	-	.4	.3	3.3	2.4	-	-	
<i>Thuja occidentalis</i> L.	-	-	-	-	-	-	-	-	-	-	-	34.0	16.8	160.2	80.4	-	-	
Stratum total	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

(Table 16 continued on next page)

(Table 16,--continued)

Tree seedling	Isabella Plantations												BWCAW				
	11-15 n = 3			16-20 n = 3			21-25 n = 1			NATURE n = 19							
	Foliage \bar{X}	SE	Foliage \bar{X}	SE	Foliage \bar{X}	SE	Foliage \bar{X}	SE	Foliage \bar{X}	SE	Foliage \bar{X}	SE					
<i>Abies balsamea</i> (L.) Mill.	0.2	-	1.8	-	40.8	40.3	346.2	342.1	4.4	9.4	55.2	13.8	119.5	30.2			
<i>Acer rubrum</i> L.	.7	0.3	2.7	1.5	.6	-	8.5	-			1.6	.4	11.9	9.1			
<i>Betula lapyrifera</i> Marsh.											1.5	1.5	38.6	17.0			
<i>Fraxinus nigra</i> Marsh.											.3	.2	1.3	.9			
<i>Picea glauca</i> (Moench) Voss.											2.3	.8	3.8	1.4			
<i>Picea mariana</i> (Mill.) B.S.P.											20.5	.6	36.7	16.1			
<i>Pinus resinosa</i> Ait.											1.1	.2	2.1	1.9			
<i>Pinus strobus</i> L.											4.9	1.8	8.3	5.6			
<i>Populus tremuloides</i> Michx.	48.4	8.1	340.0	46.1	36.4	25.8	280.5	196.6			.1	-	.4	-			
<i>Quercus rubra</i> L.											2.4	.5	12.8	2.7			
<i>Sorbus americana</i> Marsh.											35.0	33.6	76.8	74.3			
<i>Thuja occidentalis</i> L.																	
<i>Tilia americana</i> L.																	
Stratum total	49.3		344.5		77.8		635.2		4.4		124.2		340.3				
Tall shrub																	
<i>Acer spicatum</i> Lam.																	
<i>Athus crispia</i> (Ait.) Pursh.	11.3	11.0	69.7	68.2	16.6	8.5	120.0	61.4		110.9	1,057.1		9.5	4.0	56.6	27.9	
<i>Ameancher</i> Medic. spp.	10.3	9.3	45.6	41.4	9.5	-	39.5	-		29.6		4.1	4.6	68.4	32.8		
<i>Cornus rugosa</i> Lam.	42.2	13.4	371.7	114.6	53.3	21.8	475.5	195.4		121.3	1,054.3		1.0	3.3	48.3	37.4	
<i>Corylus cornuta</i> Marsh.																	
<i>Lonicera canadensis</i> Marsh.																	
<i>Lonicera hirsuta</i> Et.																	
<i>Lonicera oblongifolia</i> (Goldie) Hook.	2.9	-	11.6	-	.1	-	.2	-			<.1		<.1				
<i>Lonicera villosa</i> (Michx.) R.S.	11.4	10.9	101.3	94.8	13.3	12.4	139.4	126.9			.1		.9				
<i>Prunus pensylvanica</i> L.F.	2.4	-	12.0	-	7.2	-	37.2	-			.1		1.3				
<i>Prunus virginiana</i> L.																	
<i>Salix bebbiana</i> Sarg.	10.9	-	83.2	-	70.9	47.7	790.4	595.7			5.5	2.5	74.8	20.1			
<i>Salix</i> L. spp.																	
<i>Viburnum rafinesqueianum</i> Schult.																	
Stratum total	91.4		695.1		171.6		1,606.9		261.8		2,111.4		65.6		490.0		

(Table 16 continued on next page)

Table 16. --(continued)

(Table 16 continued on next page)

Table 16. --(continued)

Table 18 contains all new pages

(Table 16.--continued)

	Isabellia Plantations						BNCAW								
	11-15 n = 3			16-20 n = 3			Total			21-25 n = 1			Mature n = 19		
	Foliage	Total	X	Foliage	Total	X	Foliage	Total	X	Foliage	Total	X	Foliage	Total	X
<hr/>															
Fern and fern allies															
<i>Athyrium filix-femina</i> (L.) Roth.													<0.1		
<i>Dryopteris spinulosa</i> (O.F. Muell.) Watt													.3	0.2	
<i>Equisetum</i> L. spp.													.1	-	
<i>Gymnocarpium dryopteris</i> (L.) Neimn.													.9	.6	
<i>Lycopodium annotinum</i> L.													5.1	4.5	
<i>Lycopodium clavatum</i> L.													2.4	1.1	
<i>Lycopodium complanatum</i> L.													.6	.4	
<i>Lycopodium obscurum</i> L.													1.8	1.0	
<i>Polyodium virginianum</i> L.													.1	-	
<i>Pteridium aquilinum</i> (L.) Kuhn.													11.5	6.8	
Stratum total			288.1	61.8		82.9	42.4						22.9		
Moss													1,310.0	182.9	
<i>Calliergonella schreberi</i> (Willd., Br. Sch.) Grout															
<i>Dicranum</i> Hedw. spp.													232.9	58.0	
<i>Hylocladia splendens</i> (Hedw.) B.S.G.													34.8	25.6	
<i>Hypnum cristatae-castreum</i> Hedw.													26.1	10.0	
<i>Polytrichum</i> Hedw. spp.													46.4	20.5	
All other mosses													257.2	38.0	
Stratum total													1,907.4		
Lichen															
<i>Cladonia mitis</i> Sandst.													133.1	49.3	
<i>Cladonia rangiferina</i> (L.) Web.													52.1	27.2	
<i>Cladonia sylvatica</i> (L.) Hoffm.													.2	-	
<i>Peltigera aphthosa</i> (L.) Willd.													30.1	7.9	
All other lichens															
Stratum total													215.5		
Community total													62,782.4	54,781.8	
													141,313.2		

Table 17.--Aboveground oven dry biomass by species and stratum for the white-cedar community type of northeastern Minnesota

Stratum/species	(In Kg/ha)				
			BWC AW MATURE (n = 10)		Total
	Foliage		X	SE	
Tree					
Abies balsamea (L.) Mill.	2,863.5	736.2	22,121.0	5,776.6	
Betula papyrifera Marsh.	3,756.8	389.5	73,613.1	7,074.5	
Picea glauca (Moench) Voss.	707.4	220.0	7,038.8	2,273.8	
Picea mariana (Mill.) B.S.P.	102.9	77.5	975.0	732.4	
Pinus strobus L.	355.8	327.1	7,821.7	7,113.4	
Populus tremuloides Michx.	325.8	283.6	17,961.8	15,940.9	
Sorbus americana Marsh.	8.1	5.7	91.9	64.8	
Thuja occidentalis L.	6,231.5	1,023.2	69,034.2	11,714.3	
Stratum total	14,351.8		198,657.5		
Sapling					
Abies balsamea (L.) Mill.	665.1	318.5	5,616.6	2,677.3	
Betula papyrifera Marsh.	52.4	27.8	1,875.1	974.2	
Picea glauca (Moench) Voss.	2.2	1.6	14.9	11.2	
Picea mariana (Mill.) B.S.P.	34.4	24.8	212.7	156.7	
Pinus strobus L.	2.3	-	40.0	-	
Populus tremuloides Michx.	18.1	-	455.9	-	
Sorbus americana Marsh.	1.1	.5	11.6	4.8	
Thuja occidentalis L.	822.0	337.9	3,964.5	1,640.3	
Stratum total	1,597.6		12,191.3		
Tree seedling					
Abies balsamea (L.) Mill.	48.8	10.6	100.8	22.6	
Betula papyrifera Marsh.	.7	.2	13.2	3.3	
Fraxinus nigra Marsh.	.1	-	.3	-	
Picea glauca (Moench) Voss.	2.3	.6	3.7	1.1	
Picea mariana (Mill.) B.S.P.	1.8	1.4	3.1	2.5	
Pinus strobus L.	.2	.1	1.2	.9	
Populus tremuloides Michx.	.2	.2	1.1	-	
Sorbus americana Marsh.	10.1	3.7	56.8	21.4	
Thuja occidentalis L.	85.0	36.8	163.8	81.4	
Stratum total	149.2		344.0		
Tall shrub					
Acer spicatum Lam.	206.9	45.3	2,166.0	573.1	
Alnus crispa (Ait.) Pursh.	<.1	-	<.1	-	
Amelanchier Medic. spp.	<.1	<.1	<.1	<.1	
Cornus rugosa Lam.	<.1	-	.1	-	
Corylus cornuta Marsh.	36.0	17.4	244.9	124.8	
Lonicera canadensis Marsh.	2.1	.4	7.1	1.2	
Lonicera villosa (Michx.) R.D.S.	.1	-	.1	-	
Prunus virginiana L.	<.1	<.1	<.1	<.1	
Taxus canadensis Marsh.	12.8	8.8	23.2	16.0	
Stratum total	258.0		2,441.5		

(Table 17 continued on next page)

(Table 17.-continued)

Stratum/species	BWCA MATURE (n = 10)		Total	
	Foliage \bar{X}	SE	Foliage \bar{X}	SE
Low shrub				
<u>Chimaphila umbellata</u> (L.) Bart.	0.2	0.1		
<u>Diervilla Lonicera</u> Mill.	2.5	2.2		
<u>Gaultheria hispida</u> (L.) Muhl.	.3	.2		
<u>Juniperus communis</u> L.	.4	.3		
<u>Ribes americanum</u> Mill.	<.1	-		
<u>Ribes glandulosum</u> Grauer.	<.1	-		
<u>Ribes lacustre</u> (Pers.) Poir.	<.1	-		
<u>Ribes triste</u> Pall.	<.1	-		
<u>Rubus parviflorus</u> Nutt.	11.4	6.6		
<u>Rubus pubescens</u> Raf.	3.6	1.8		
<u>Rubus strigosus</u> Michx.	.8	.7		
Stratum total	19.3			
Herb				
<u>Aralia nudicaulis</u> L.	17.1	4.3		
<u>Aster macrophyllus</u> L.	10.7	3.5		
<u>Carex</u> L. spp.	7.6	3.7		
<u>Circaeа alpina</u> L.	.6	.5		
<u>Clematis virginiana</u> L.	.1	-		
<u>Clintonia borealis</u> (Ait.) Raf.	9.3	2.2		
<u>Coptis groenlandica</u> (Oeder) Fern.	1.2	.4		
<u>Cornus canadensis</u> L.	11.3	2.3		
<u>Galium triflorum</u> Michx.	.7	.4		
<u>Goodyera repens</u> (L.) R. Br.	.3	.2		
Grasses	4.1	2.2		
<u>Habenaria orbiculata</u> (Pursh) Torr.	.5	-		
<u>Impatiens capensis</u> Meerb.	3.8	2.9		
<u>Lathyrus ochroleucus</u> Hook.	.2	-		
<u>Linnaea borealis</u> L.	6.1	3.2		
<u>Maianthemum canadense</u> Desf.	4.2	1.5		
<u>Mertensia paniculata</u> (Ait.) G. Don.	.7	-		
<u>Mitella nuda</u> L.	16.1	4.0		
<u>Moneses uniflora</u> (L.) Gray	.1	.1		
<u>Oxalis montana</u> Raf.	.5	.5		
<u>Pyrola secunda</u> L.	.6	.2		
<u>Pyrola virens</u> Schweigg.	.2	.2		
<u>Streptopus roseus</u> Michx.	2.7	1.6		
<u>Trifoliate borealis</u> Raf.	2.1	1.1		
<u>Viola</u> L. spp.	1.8	.6		
Stratum total	102.6			
Fern and fern allies				
<u>Athyrium filix-femina</u> (L.) Roth.	0.7	-		
<u>Botrychium virginianum</u> (L.) Sw.	.6	-		
<u>Cystopteris fragilis</u> (L.) Bernh.	.1	-		
<u>Dryopteris phegopteris</u> L.	.1	-		
<u>Dryopteris spinulosa</u> (O.F. Muell.) Watt	11.0	4.7		
<u>Gymnocarpium trypteris</u> (L.) Newm.	5.4	2.3		
<u>Lycopodium annotinum</u> L.	1.9	1.0		
<u>Lycopodium clavatum</u> L.	1.6	.8		
<u>Lycopodium lucidulum</u> Michx.	1.3	.8		
<u>Lycopodium obscurum</u> L.	1.9	1.6		
Stratum total	24.6			
Moss				
<u>Calliergonella schreberi</u> (Willd., BR. & Sch.) Grout	100.4	57.4		
<u>Dicranum</u> Hedw. spp.	8.5	4.7		
<u>Hylocomium splendens</u> (Hedw.) B.S.G.	44.3	19.5		
<u>Hypnum crista-castrensis</u> Hedw.	1.3	.5		
<u>Polytrichum</u> Hedw. spp.	5.4	4.3		
All other mosses	533.3	89.6		
Stratum total	693.2			
Lichen				
<u>Cladonia rangiferina</u> (L.) Web.	0.5	-		
All other lichens	16.9	7.7		
Stratum total	17.4			
Community total	214,491.4			

Ohmann, Lewis F.; Grigal, David F.

Plant species biomass estimates for 13 upland community types of northeastern Minnesota. Resour. Bull. NC-88. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1985. 52 p.

Presents total aboveground oven-dry biomass estimates by stratum and species for 13 plant community types located in northeastern Minnesota.

KEY WORDS: Plantation, biomass estimation, dry weight, silviculture, plant community.